



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

The biochemistry and molecular biology of chlorophyll breakdown

Kuai, Benke ; Chen, Junyi ; Hörtensteiner, Stefan

Abstract: Chlorophyll breakdown is one of the most obvious signs of leaf senescence and fruit ripening. The resulting yellowing of leaves can be observed every autumn, and the color change of fruits indicates their ripening state. During these processes, chlorophyll is broken down in a multistep pathway, now termed the 'PAO/phylobilin' pathway, acknowledging the core enzymatic breakdown step catalysed by pheophorbide a oxygenase, which determines the basic linear tetrapyrrole structure of the products of breakdown that are now called 'phylobilins'. This review provides an update on the PAO/phylobilin pathway, and focuses on recent biochemical and molecular progress in understanding phylobilin-modifying reactions as the basis for phylobilin diversity, on the evolutionary diversity of the pathway, and on the transcriptional regulation of the pathway genes.

DOI: <https://doi.org/10.1093/jxb/erx322>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-148880>

Journal Article

Accepted Version

Originally published at:

Kuai, Benke; Chen, Junyi; Hörtensteiner, Stefan (2017). The biochemistry and molecular biology of chlorophyll breakdown. *Journal of Experimental Botany*, 69(4):751-767.

DOI: <https://doi.org/10.1093/jxb/erx322>

The Biochemistry and Molecular Biology of Chlorophyll Breakdown

Benke Kuai^{1,2}, Junyi Chen^{1,2} & Stefan Hörtensteiner^{3*}

¹State Key Laboratory of Genetic Engineering, School of Life Sciences, Fudan University, Shanghai 200438, China

²Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, Institute of Biodiversity Science, Fudan University, Shanghai 200438, China

³Institute of Plant and Microbial Biology, University of Zurich, Zollikerstrasse 107, CH-8008 Zurich, Switzerland

Email Benke Kuai: bkkuai@fudan.edu.cn

Email Junyi Chen: chenjy14@fudan.edu.cn

*To whom correspondence should be addressed.

E-mail: shorten@botinst.uzh.ch

Tel: +41 44 634 82 82

Fax: +41 44 634 82 04

Date of re-submission: August 10, 2017

Number of Figures: 2 (both in color in print); number of Tables: 1

Total word count: 15373

Running title: Chlorophyll breakdown

27 **Abstract**

28 Chlorophyll breakdown is one of the most obvious signs of leaf senescence and fruit ripening.
29 The resulting yellowing of leaves can be observed every autumn, and the color change of
30 fruits indicates their ripening state. During these processes, chlorophyll is broken down in a
31 multistep pathway, now termed ‘PAO/phyllobilin’ pathway, acknowledging the core
32 enzymatic breakdown step catalyzed by pheophorbide *a* oxygenase. It determines the basic
33 linear tetrapyrrole structure of the products of breakdown that are now called ‘phyllobilins’.
34 This review provides an update on the PAO/phyllobilin pathway, thereby focusing on recent
35 biochemical and molecular progresses in understanding phyllobilin modifying reactions as the
36 basis for phyllobilin diversity, on evolutionary diversity of the pathway, and on transcriptional
37 regulation of pathway genes.

38

39

40 **Introduction**

41 During green organ senescence/maturation as well as during fruit ripening, efficient
42 chlorophyll (Chl) breakdown causes rapid degreening, and consequently unmasks the
43 existence of yellow carotenoids and splendidly highlights the synthesis of diverse anthocyanins
44 during the late growing season. Coordinated Chl breakdown is an integral process of
45 senescence/maturation that is developmentally programmed to facilitate the dynamic
46 remobilization of nutrients from senescent organs/tissues to growing organs,
47 reproductive/storage organs in particular (Lim *et al.*, 2007). It has been shown that
48 chloroplasts contain up to 75% of the nitrogen in photosynthetic tissues, with Rubisco
49 accounting for 20-30% of total leaf nitrogen (Makino and Osmond, 1991; Peoples and Dalling,
50 1988), and up to 95% of seed proteins is synthesized from amino acids derived from the
51 degradation of existing proteins in leaves (Taylor *et al.*, 2010). It is vital, on one hand, to
52 abolish the phototoxicity of free Chl molecules that are uncoupled from light-harvesting Chl-
53 binding complex proteins (LHCs) of the thylakoids in order to maintain cell viability to allow
54 for an efficient remobilization of nutrients in senescent organs/tissues. On the other hand, Chl
55 breakdown has been repeatedly shown to be a prerequisite for the degradation of LHCs in
56 senescent leaves and as such important for accessing this second largest pool of chloroplast
57 nitrogen (Feller *et al.*, 2008; Hörtensteiner and Feller, 2002; Wu *et al.*, 2016). In addition, by-
58 products of Chl degradation are re-used for biosynthetic purposes (Christ and Hörtensteiner,
59 2014); particularly phytol, which is salvaged into tocopherol during leaf senescence (vom

Dorp *et al.*, 2015).

The biochemical pathway of Chl breakdown, the so-called ‘PAO/phyllobilin pathway’ (Christ and Hörtensteiner, 2014; Kräutler and Hörtensteiner, 2014), has been almost completely revealed based on the characterized enzymes responsible for catalyzing Chl degradation, particularly with the recent demonstration that NONYELLOWING (NYE) proteins, also termed STAY-GREEN (SGR) possess Mg-dechelataase activity (Shimoda *et al.*, 2016) (Figure 1). The pathway involves two distinct phases: the early phase deals with the degradation of phototoxic free Chl molecules and their intermediates at the thylakoid membrane within the chloroplast, and the late phase is responsible for modifications of colorless Chl catabolites and their translocation from the chloroplast to the vacuole. Recently, new insight has been obtained on the topology of Chl degradation within the chloroplast as well as on cytosolic modifications of colorless catabolites. Although Chl breakdown can be initiated in various tissues under diverse scenarios, e.g. upon pathogen infection (Mur *et al.*, 2010) and under abiotic stresses (Munné-Bosch and Alegre, 2004), massive Chl breakdown occurs mainly in senescent leaves. By taking advantage of the well-characterized *Chl catabolic genes (CCGs)*, the molecular regulation of Chl breakdown was explored during the past few years, and some of the transcriptional regulatory mechanisms have been revealed in *Arabidopsis thaliana* (Figure 2) as well as in other species (Chen *et al.*, 2017; Delmas *et al.*, 2013; Gao *et al.*, 2016; Li *et al.*, 2016; Liang *et al.*, 2014; Oda-Yamamizo *et al.*, 2016; Qiu *et al.*, 2015; Sakuraba *et al.*, 2016; Sakuraba *et al.*, 2014a; Song *et al.*, 2014; Yin *et al.*, 2016; Zhang *et al.*, 2015; Zhu *et al.*, 2015). This review aims to provide an updated summary of progresses in the elucidation of the PAO/phyllobilin pathway and its transcriptional regulation, and to discuss the significance of these progresses as well as remaining lack of knowledge that demands further investigations.

The PAO/phyllobilin pathway of chlorophyll breakdown

Until more than 25 years ago, the fate of Chl during senescence was enigmatic (Hendry *et al.*, 1987). Thus, the identification of compounds that likely derived from Chl and that accumulated in a senescing variety of *Festuca pratensis*, but were absent from a respective stay-green variety, marked an important breakthrough in the field (Matile *et al.*, 1987). In 1991, *Hv-NCC-1* isolated from senescent barley (*Hordeum vulgare*) leaves was structurally characterized as the first catabolite of Chl that accumulates in the vacuoles of senescing cells

and can be considered as an ultimate product of degradation (Kräutler *et al.*, 1991). This important step fostered the research on Chl breakdown and has led to the identification of currently over 40 breakdown products of Chl from more than 20 plant species that possess the basic common structure of a Chl *a*-derived linear tetrapyrrole, but differ from each other in terms of characteristic side chain modifications and/or stereoisomeric properties. Collectively these catabolites are now termed phyllobilins (Kräutler, 2014).

Chlorophyll is converted to phyllobilins

Phyllobilins can be categorized into two different groups (Kräutler, 2014, 2016). Type I phyllobilins are the historically first identified nonfluorescent Chl catabolites (NCCs) that structurally are 1-formyl, 19-oxobilins (see Figure 1, for atom and pyrrole ring numbering of phyllobilins), while the only recently discovered type II phyllobilins are 1,19-dioxobilins (Müller *et al.*, 2011), now termed dioxobilin-type nonfluorescent Chl catabolites (DNCCs). Interestingly, the later are similar to biliverdin, the first product of heme degradation, in which the ‘northern’ meso carbon is lost as CO during the heme oxygenase-catalyzed porphyrin ring opening reaction (Unno *et al.*, 2007). In NCCs and DNCCs, all three remaining methine bridges between the pyrrole rings that remain after porphyrin ring opening are fully reduced, leaving behind a linear tetrapyrrole backbone with an unconjugated electron system. The nonfluorescent NCCs and DNCCs are derived from fluorescent precursors, termed FCCs and DFCCs, respectively, that are isomerized to the nonfluorescent end products by acid-catalyzed tautomerization at pyrrole ring C, the isocyclic ring E and the ‘southern’-meso position. This isomerization was shown to be driven by an intramolecular protonation step involving the C12 propionic acid side chain (Oberhuber *et al.*, 2003) and to occur in a pH-dependent manner (Christ *et al.*, 2012) inside the acidic vacuolar sap, the final storage place of the phyllobilins (Hinder *et al.*, 1996; Matile *et al.*, 1988). As mentioned above, DNCCs and NCCs possess characteristic side chain modifications at different peripheral positions. So far, the following functionalizations have been described: (i) hydroxylation at C3² that can be followed by glycosylation and/or malonylation (R¹ in Figure 1), (ii) demethylation at O8⁴ (R²), (iii) dihydroxylation of the C18 vinyl group, that can be followed by glucosylation (R³), and (iv) hydroxymethylation at either C2 or C4 (R⁴). It was been assumed that these modifications are introduced in the fluorescent FCC and/or DFCC precursors before their import into the vacuole and subsequent isomerization to NCCs and DNCCs.

Interestingly, in a few plant species, i.e. banana (*Musa acuminata*) and *Spathiphyllum wallisii*, so-called *hypermodified* FCCs (*hmFCCs*) were identified that persist in senescent leaves or fruits and are not isomerized to the respective NCCs (Kräutler, 2014, 2016). This has been rationalized by functionalization of the C12 propionic acid with different moieties (R^5 in Figure 1) such as daucyl, glucosyl, digalactosyl-glyceryl or glucosyl-(dihydroxyphenyl) ethyl, which dramatically slows down FCC-to-NCC isomerization (Oberhuber *et al.*, 2008). Nevertheless, a few NCCs with a conjugated C12 propionic acid side chain have been described (Moser *et al.*, 2012). Among them is a bicycloglycosidic NCC from *Ulmus glabra* in which a glucose moiety intramolecularly connects C3² with C12³ (Scherl *et al.*, 2016). Apart from these type I and type II phyllobilins, small quantities of so-called yellow and pink Chl catabolites (YCCs and PiCCs, respectively) have been identified in senescent leaves from several species (Moser *et al.*, 2009b; Moser *et al.*, 2008b; Ulrich *et al.*, 2011). They are oxidation products of NCCs that have been considered to naturally occur as the result of plant endogenous oxidative activities (Vergeiner *et al.*, 2015) and that might contribute to the color of senescent leaves or ripened fruits (Moser *et al.*, 2012; Moser *et al.*, 2008b). Although so far, DNCC-derived dioxobilin-type YCCs or PiCCs have not been described (Kräutler, 2016), such oxidations are likely not restricted to NCCs.

The biochemistry and topology of chlorophyll breakdown

The biochemical mechanism of Chl breakdown was elucidated stepwise during the last 20 years. With the elucidation of the first phyllobilin structure in 1991 (Kräutler *et al.*, 1991) and inhibition studies (Hörtensteiner *et al.*, 1995), it became obvious that a Fe-dependent oxygenase is responsible for porphyrin ring cleavage, thus determining the common backbone structure of all subsequent intermediary and end products of breakdown. This oxygenase was later identified as a Rieske type monooxygenase that oxygenolytically opens the porphyrin ring of pheophorbide *a*, a Mg^{2+} - and phytol-free intermediate of Chl *a* breakdown, and was thus named PHEOPHORBIDE *a* OXYGENASE (PAO) (Pružinská *et al.*, 2003). Because of the importance of this step in the pathway, it is nowadays referred to as the ‘PAO/phyllobilin’ pathway of Chl breakdown (Kräutler and Hörtensteiner, 2014). Generally, the PAO/phyllobilin pathway can be divided into two parts. Part I was considered to include all reactions that are involved in the conversion of Chl *b* (via Chl *a*) and Chl *a* to *primary* FCC (*pFCC*) (Christ and Hörtensteiner, 2014). All of these reactions occur inside of the chloroplast and are common to all plant species that have been analyzed for Chl breakdown so far. The

recent identification of the C³-hydroxylase as a chloroplast inner envelope-located Rieske type oxygenase, named TRANSLOCON AT THE INNER CHLOROPLAST ENVELOPE 55 (TIC55) (Hauenstein *et al.*, 2016), and the common occurrence of NCCs and DNCCs that are derived from C³-hydroxylated *p*FCC (hydroxy-*p*FCC) calls for including this step in reactions of part I of the PAO/phyllobilin pathway. By contrast, part II describes all further side chain-modifying reactions, all of which take place outside the chloroplast, i.e. at the ER or in the cytosol, and include the non-enzymatic isomerization of fluorescent to nonfluorescent phyllobilins.

Reactions of part I of the PAO/phyllobilin pathway

Conversion of Chl *b* to Chl *a* – Except one (Müller *et al.*, 2006), all NCCs and DNCCs that are known up to now derive from Chl *a*. The reason for this intriguing fact is the specificity of two of the downstream Chl catabolic enzymes (CCEs), i.e. the Mg-dechelateses, termed SGR or NYE (the latter name being used in the following) and PAO, for Chl *a* and pheophorbide *a*, respectively, while respective '*b*' pigments are not accepted as substrates (Hörtensteiner *et al.*, 1995; Shimoda *et al.*, 2016). Thus, prior to its degradation by the PAO/phyllobilin pathway, Chl *b* must be converted to Chl *a* (Shimoda *et al.*, 2012). This is achieved by the reductive half of the so-called 'chlorophyll cycle', which interconverts chlorophyll(ide) *a* and chlorophyll(ide) *b* (Tanaka and Tanaka, 2011). Chl *b* to Chl *a* reduction is a two-step reaction catalyzed by NONYELLOW COLORING 1 (NYC1) and NYC1-LIKE (NOL), two paralogous Chl *b* reductases that depend on NADPH, and HYDROXYMETHYL CHL *a* REDUCTASE (HCAR), which depends on reduced ferredoxin (Fd) as a source of electrons. Genes encoding respective CCEs were recently identified in rice (*Oryza sativa*) and Arabidopsis (Horie *et al.*, 2009; Kusaba *et al.*, 2007; Meguro *et al.*, 2011; Sato *et al.*, 2009).

Mg-dechelation and dephytylation – As mentioned above, pheophorbide *a* is the substrate of PAO, indicating that Mg-dechelation and dephytylation happen before porphyrin ring cleavage. Since its identification more than a century ago (Willstätter and Stoll, 1913), CHLOROPHYLLASE (CLH) that hydrolyzes Chl to chlorophyllide and phytol was assumed to be involved in Chl degradation during leaf senescence. In favor of this are high *in vitro* activities of CLHs (Arkus *et al.*, 2005; Schelbert *et al.*, 2009) and their wide distribution in higher plants and algae (Drazkiewicz, 1994; Shioi and Sasa, 1986). Since 1999, when the first CLHs were cloned independently by two groups (Jakob-Wilk *et al.*, 1999; Tsuchiya *et al.*,

1999), *CLH* genes have been identified from many plant species with surprising results: only in a few cases were the cloned proteins predicted or experimentally shown to be targeted to the chloroplast (Azoulay-Shemer *et al.*, 2011; Azoulay Shemer *et al.*, 2008), while for example the two Arabidopsis CLHs localize outside the plastid, i.e. in the ER and tonoplast (Hu *et al.*, 2015; Schenk *et al.*, 2007). Originally, this was interpreted as Chl dephytylation to (additionally) occur outside the plastid (Takamiya *et al.*, 2000); however, the involvement of CLHs in leaf senescent-related Chl breakdown was repeatedly challenged (Hörtensteiner, 2006; Hu *et al.*, 2015; Liao *et al.*, 2007; Schelbert *et al.*, 2009; Schenk *et al.*, 2007; Zhou *et al.*, 2007). Recently, CLH (together with Chl) was proposed to play a role as a plant endogenous two component defense system. In this system, CLH gets access to plastid-localized Chl upon cell damage by leaf-chewing herbivores, causing the production of chlorophyllide, which was shown to hamper development of *Spodoptora litura* larvae (Hu *et al.*, 2015).

Using an *in silico prediction* approach, Arabidopsis candidates for a phytol hydrolase that is chloroplast-localized and senescence-regulated were proposed (Schelbert *et al.*, 2009). One of them, now named PHEOPHYTINASE (PPH), was shown to specifically hydrolyze pheophytin and respective mutants were unable to degrade Chl and thus exhibited a stay-green phenotype. PPH was independently identified by two other groups in Arabidopsis (Ren *et al.*, 2010) and rice (Morita *et al.*, 2009). Since then PPH has been characterized in many other plant species and consistently shown to be required for Chl breakdown during leaf senescence (Büchert *et al.*, 2011; Cheng and Guan, 2014; Guyer *et al.*, 2014; Zhang *et al.*, 2016). This solidified the concept that Mg-dechelation precedes dephytylation in the PAO/phyllobilin pathway (Kusaba *et al.*, 2013; Tanaka *et al.*, 2011).

Many activities have been considered in the past to be responsible for Mg-dechelation. These include (i) non-enzymatic loss of the Mg²⁺-ion as the result of changes in plastidial pH during senescence – Mg is known to be readily lost from Chl under acidic conditions (Saga and Tamiaki, 2012), (ii) involvement of potential Mg-chelators, i.e. heat-stable, low molecular weight compounds termed Mg-chelating substances (Costa *et al.*, 2002; Suzuki and Shioi, 2002), and (iii) Mg-dechelatasases (Suzuki and Shioi, 2002; Vicentini *et al.*, 1995). However, the molecular nature of such dechelatasases remained enigmatic until recently, because of the unawareness that Chl is the likely substrate for dechelation (Schelbert *et al.*, 2009; Shimoda *et al.*, 2016), not the much more polar chlorophyllide or the often used artificial substrate chlorophyllin (Suzuki and Shioi, 2002; Vicentini *et al.*, 1995). Only very recently, NYE was shown to function as Mg-dechelataase *in vitro* as well as *in vivo* (Shimoda *et al.*, 2016). NYE

has been identified independently by several groups (Armstead *et al.*, 2006, 2007; Jiang *et al.*, 2007; Park *et al.*, 2007; Ren *et al.*, 2007), but mutants that are now known to be deficient in NYE exist since decades or even centuries (Barry, 2009; Hörtensteiner, 2009). Indeed, the green cotyledon variety of pea (*Pisum sativum*) used by Gregor Mendel for the dissection of the laws of genetics in 1866 (Mendel, 1866), has a defect in the *NYE1* gene (Aubry *et al.*, 2008; Sato *et al.*, 2007). In the genome of Arabidopsis, three NYEs are encoded, i.e. the highly homologous NYE1 and NYE2 proteins and NYE/SGR-like (SGRL), which is more distantly related. A major difference of these proteins is the presence in NYE1 and NYE2 of a highly conserved cysteine-rich C-terminal domain of unknown function that is absent in SGRL (Hörtensteiner, 2009). It may explain their distinct substrate specificities: while NYE1 is highly specific-for Chl *a*, SGRL prefers phytol-free chlorophyllide *a* as substrate (Shimoda *et al.*, 2016). Different roles of the three Arabidopsis NYEs have been proposed in the past, some of which in retrospect and in light of their role as Mg-dechelatasers may be questioned. Firstly, NYE1 was proposed as a mere scaffold protein whose function is to recruit other CCEs to form a highly dynamic protein complex at LHCII, to allow metabolite channeling of potentially phototoxic Chl breakdown intermediates (Sakuraba *et al.*, 2012). As Mg-dechelatare and first enzyme in Chl *a* degradation, NYE1 still could exert this function to coordinate metabolism downstream of Chl *a*, however its demonstrated interaction with upstream enzymes such as NYC1 or NOL is difficult to rationalize. Secondly, NYE2 has been proposed to have a negative role in Chl breakdown (Sakuraba *et al.*, 2014c), but in retrospect it seems not reasonable why an Mg-dechelatare should act negatively. It has to be noted, however, that this proposed negative role of NYE2 was refuted recently (Wu *et al.*, 2016). Thirdly, SGRL, whose expression is high in seedlings but declines during senescence, was proposed to have a role in stress-related Chl degradation rather than for leaf senescence (Sakuraba *et al.*, 2014b). Its high specificity for chlorophyllide *a* (Shimoda *et al.*, 2016) would require SGRL to function together with a CLH-like enzyme rather than with PPH such as the recently identified CHLOROPHYLL DEPHYTYLASE 1 (Lin *et al.*, 2016). Nevertheless, SGRL was shown to physically interact with PPH (Sakuraba *et al.*, 2014b). In summary, more research is required to determine the exact functions of these NYEs for Mg-dechelation of Chl.

Formation of *p*FCC from pheophorbide *a* – The conversion of pheophorbide *a* to *p*FCC was demonstrated to be a Fd-dependent two-step reaction, because besides Fd, two additional protein fractions were shown to be required for *in vitro* *p*FCC formation (Hörtensteiner *et al.*,

1995; Rodoni *et al.*, 1997), i.e. a membrane fraction of senescent chloroplasts later shown to be PAO (Pružinská *et al.*, 2003), and a stromal fraction, later identified as red Chl catabolite reductase (RCCR) (Wüthrich *et al.*, 2000). The intermediary product of the PAO reaction, red Chl catabolite (RCC), is metabolically channeled *in vitro*, explaining the simultaneous requirement in respective assays of both enzymes. In line with this, PAO and RCCR were shown to physically interact (Pružinská *et al.*, 2007; Sakuraba *et al.*, 2012).

PAO is a Rieske type monooxygenase with two C-terminally located transmembrane domains that anchors it within the thylakoid membrane (Sakuraba *et al.*, 2012). Using $^{18}\text{O}_2$ labeling experiments it was demonstrated that only the oxygen incorporated at the C1 formyl position of *pFCC* is derived from molecular oxygen (Hörtensteiner *et al.*, 1998). Intriguingly, similar experiments with the green alga *Auxenochlorella protothecoides* that efficiently degrades Chl under N-limiting, heterotrophic conditions in the dark (Engel *et al.*, 1996), uncovered the principle same type of mechanism (Curty *et al.*, 1995). In contrast to higher plants, *A. protothecoides* does not further convert RCC, but excretes it into the surrounding medium. Accordingly, RCCR is absent from this green alga as based on recent genome and RNAseq analysis (Gao *et al.*, 2014) (Aubry & Hörtensteiner, unpublished).

RCCR specifically reduces the C15/C16 double bond of RCC, thereby introducing a new stereo center at C16. This reduction occurs in a highly stereospecific manner and was shown to depend on the source of RCCR (Hörtensteiner *et al.*, 2000b; Pružinská *et al.*, 2007), dividing these proteins into type I or type II RCCR. For example the Arabidopsis enzyme (a type I RCCR) produces specifically *pFCC* (Mühlecker *et al.*, 1997), the presumed ‘S’ stereoisomer (Oberhuber *et al.*, 2008), while the enzymes from bell pepper (*Capsicum annuum*) or tomato (*Solanum lycopersicum*) produce *epi-pFCC* (Mühlecker *et al.*, 2000), i.e. the presumed C16 ‘R’ epimer (Oberhuber *et al.*, 2008). Analysis of chimeric RCCRs composed of Arabidopsis and tomato sequence parts identified a single amino acid residue, which when exchanged with the respective tomato residue, converted Arabidopsis RCCR to a type II enzyme that produces *epi-pFCC* (Pružinská *et al.*, 2007). The stereochemistry introduced by RCCR is reflected in the final phyllobilins, i.e. each of the above described NCCs and DNCCs may, depending on the plant species occur in two C16-isomeric forms. RCCRs are members of the Fd-dependent bilin reductase (FDBR) family that includes ELONGATED HYPOCOTYL 2, i.e. phytochromobilin synthase of higher plants and several bilin reductases of cyanobacteria and algae that catalyze phycobilin biosynthesis (Frankenberg and Lagarias, 2003). Although the 3D structures of RCCR and other FDBRs are

highly similar (Sugishima *et al.*, 2009; Sugishima *et al.*, 2010), i.e. pointing to a similar radical mechanism of electron transfer in RCCR as shown for some other FDBRs (Tu *et al.*, 2004), RCCR exhibits very low sequence similarity with these other FDBRs (Frankenberg *et al.*, 2001).

Mutants in PAO and RCCR were originally identified as ACCELERATED CELL DEATH (ACD) 1 and ACD2, respectively (Greenberg and Ausubel, 1993; Greenberg *et al.*, 1994). PAO and RCCR mutants exhibit a light-dependent cell death phenotype, which was shown to be caused by the accumulation of, respectively, pheophorbide *a* and RCC that are highly photodynamic (Mach *et al.*, 2001; Pružinská *et al.*, 2007; Pružinská *et al.*, 2003; Tanaka *et al.*, 2003; Xodo *et al.*, 2012). Interestingly, however, absence of PAO, but not RCCR (Aubry & Hörtensteiner, unpublished), causes additional light-independent cell death, indicating that besides its high phototoxicity (Xodo *et al.*, 2012) pheophorbide *a* could act as a cell death signaling molecule in the dark (Hirashima *et al.*, 2009). However, pheophorbide *a* unlikely functions as a retrograde signal itself, because it was shown to be unable to leave the plastid (Christ *et al.*, 2012), and other components of this proposed signaling pathway have not yet been identified. Once converted to *pFCC*, the potential phototoxicity of chlorophyll is abolished, and although *pFCC* still has a certain capacity to produce singlet oxygen (Jockusch *et al.*, 2014), this is far below the phototoxicity of pheophorbide *a* or RCC. Thus, formation of *pFCC* by PAO and RCCR can be considered the key step of breakdown that accomplishes the abolishment of Chl phototoxicity as the main purpose for its degradation.

***pFCC*-hydroxylation by TIC55** – All so far described reactions of the first part of the PAO/phyllobilin pathway occur in plastids. The recent identification of TIC55 as C3² hydroxylase (Hauenstein *et al.*, 2016) adds this reaction to part I, because (i) it commonly occurs in all plant species, as concluded from the presence of C3²-hydroxylated phyllobilins in all investigated species (Hauenstein *et al.*, 2016; Kräutler, 2016), and (ii) the reaction takes place in the plastid as shown by the presence in senescent Arabidopsis chloroplasts of both *pFCC* and hydroxy-*pFCC* (Hauenstein *et al.*, 2016). At first glance, this finding was rather surprising, because TIC55 was identified as a potential component of the protein import machinery at the chloroplast inner envelope (Balsera *et al.*, 2010; Calibe *et al.*, 1997); however, this had been questioned in the past (Boij *et al.*, 2009). Evidences for the role of TIC55 as *pFCC*-hydroxylase came from *in vitro* analyses that showed the requirement for reduced Fd for hydroxylating activity and from the analysis of *tic55* mutants which

exclusively accumulate non-hydroxylated phyllobilins, while in wild type more than 50% of all phyllobilins are hydroxylated (Hauenstein *et al.*, 2016).

Reactions of part II of the PAO/phyllobilin pathway

As mentioned above, over 40 structurally different phyllobilins have been isolated from more than 20 plant species. In the past, phyllobilins were named according to their source, by their type (RCC, FCC, DFCC, NCC, DNCC) and their relative or absolute retention time on a standardized HPLC system (Ginsburg and Matile, 1993; Kräutler, 2014, 2016). However, it turns out that, with this system, constitutionally identical phyllobilins receive different names (see Table 1). Thus, a simplifying nomenclature was proposed recently that classifies phyllobilins according to their type and their monoisotopic mass (Christ *et al.*, 2016). Since LC-MS analysis started to become more widely used for phyllobilin analysis (Christ *et al.*, 2016; Müller *et al.*, 2014; Rios *et al.*, 2015), this seems reasonable. Because of distinct and characteristic fragmentation patterns in tandem MS, different classes of phyllobilins such as NCCs vs DNCCs (Christ *et al.*, 2016) and fluorescent vs nonfluorescent phyllobilins (Hauenstein *et al.*, 2016) can be distinguished. However, this classification system ignores isomeric differences between otherwise identical phyllobilins, as for example C16-epimers. Nevertheless, in the past, structurally identical phyllobilins isolated from different species have obtained different names (see Table 1). Thus, it may be desirable in the future to establish a simple but conclusive nomenclature for phyllobilins that concisely describes their structural features but is not redundant.

The basis of the structural variability of phyllobilins are functionalizations at different side chains that occur species-specifically. Besides the TIC55-catalyzed C3² hydroxylation, the enzymes of only two further modifications have molecularly been identified.

C1 deformylation – Since their first description in 1991 (Kräutler *et al.*, 1991), NCCs were identified from many plant species and for a long time were considered the exclusive degradation products of Chl. Only a few isolated reports hinted to the occurrence of dioxobilin type forms of phyllobilins, originally named urobilinogenoidic Chl catabolites (Djapic and Pavlovic, 2008; Djapic *et al.*, 2009b; Losey and Engel, 2001). They were considered to be oxidation products of NCC precursors that are, thus, likely formed inside the vacuole (Losey and Engel, 2001). Occurrence of DNCCs and NCCs varies among plant species. For example, *Acer platanoides* exclusively accumulates one single DNCC (Müller *et*

al., 2011) while *Cercidiphyllum japonicum* exclusively accumulates NCCs (Curty and Engel, 1996; Oberhuber *et al.*, 2003). By contrast, *Arabidopsis* for example forms both NCCs and DNCCs simultaneously, whereby the latter account for over 90% of all phyllobilins found (Christ *et al.*, 2013). This hinted to a dedicated, species-specific activity responsible for DNCC formation. Using a combination of inhibition studies and *in silico* analysis, candidates for such an activity were identified in *Arabidopsis* (Christ *et al.*, 2013). Indeed, DNCC formation was efficiently inhibited by CO, indicating the involvement of cytochrome P450 monooxygenases (CYPs). Based on mutant analysis and *in vitro* activity determination, CYP89A9 was identified as the sole CYP enzyme responsible for DNCC formation in *Arabidopsis*. As known from many other CYPs (Schuler *et al.*, 2006), CYP89A9 localizes to the ER, and it catalyzes an oxidative deformylation of FCCs to respective DFCCs (Christ *et al.*, 2013). Intriguingly, CYP89A9 seems to be the sole member of the seven-membered CYP89 subfamily in *Arabidopsis*, because *cyp89a9* mutants are devoid of DNCCs, but accumulate accordingly more NCCs. Since the mutants do not show any obvious phenotype, phyllobilin deformylation does not seem essential. Thus, it will be interesting to analyze in the future how and why during evolution CYP89A9 was specifically recruited for phyllobilin deformylation in *Arabidopsis* and whether distinct species that accumulate DNCCs evolved this activity independently from each other.

O8⁴ demethylation – Similar to phyllobilin deformylation, demethylation of the C8²-carboxymethyl group occurs in a species-specific manner. In *Arabidopsis*, it is catalyzed by the cytosol-located METHYLESTERASE 16 (MES16) (Christ *et al.*, 2012), a member of a 20-membered protein family (Yang *et al.*, 2008), indicating its specific recruitment for Chl breakdown. Based on *in vitro* assays, demethylation has originally been considered to occur at the level of pheophorbide *a*, and be followed by spontaneous C13²-decarboxylation which leads to the formation of pyropheophorbide *a* (Shioi *et al.*, 1996; Suzuki *et al.*, 2006). However, C8²-decarboxylated phyllobilins have not been identified up to now and comparison of MES16 activity with pheophorbide *a* and *p*FCC demonstrated the specific stability of the remaining carboxylgroup in the latter (Christ *et al.*, 2012). In addition, in an *Arabidopsis paol* mutant in which MES16 is mis-targeted to the chloroplast, pyropheophorbide *a* accumulates (Christ *et al.*, 2012). This indicates that the natural substrate of MES16 are cytosolic FCCs and DFCCs. Recently, the crystal structure was resolved at high resolution, indicating that MES16 belongs to the α/β -hydrolase protein superfamily (Li

and Pu, 2016). Besides phyllobilins, MES16 was shown to specifically catalyze the demethylation of methyl-indole acetic acid, but not of other phytohormone methylesters (Christ *et al.*, 2012; Li and Pu, 2016); however, the biological relevance of this MES16 activity remains unknown.

Phyllobilin transport and isomerization – Even before the first structure identification of a phyllobilin, it was proposed that Chl catabolites are stored in the vacuole of senescing barley cells (Matile *et al.*, 1988) and this was confirmed in other plant species (Christ *et al.*, 2012; Hinder *et al.*, 1996; Matile, 1997; Pružinská *et al.*, 2007). In addition and as mentioned above, isomerization of fluorescent precursors to respective NCCs and DNCCs is driven by an acidic pH (Oberhuber *et al.*, 2003), i.e. likely happens inside the vacuole. Finally, transport of phyllobilins into isolated barley vacuoles was demonstrated to be ATP-dependent (Hinder *et al.*, 1996). Although the ATP-binding cassette (ABC) transporters ABCC2 and ABCC3 of *Arabidopsis* were shown to transport NCCs after expression in yeast (*Saccharomyces cerevisiae*) (Lu *et al.*, 1998; Tommasini *et al.*, 1998), the *in vivo* transporter(s) of phyllobilins at the tonoplast have not been identified. Likewise, export of FCCs from the chloroplast was considered to be ATP-dependent (Matile *et al.*, 1992), but respective transporters have not yet been identified.

How conserved is the PAO/phyllobilin pathway in different tissues?

Visible Chl breakdown, i.e. net loss of green color, does not only occur during natural leaf senescence, but, for example, also in response to different biotic and abiotic stresses. This is seen dramatically during the desiccation of resurrection plants that are capable to fully re-hydrate after the re-supply of water (Tuba *et al.*, 1996). In addition to these leaf-related processes, Chl breakdown also occurs in other tissues of a plant, such as during fruit ripening and seed maturation (Delmas *et al.*, 2013; Moser *et al.*, 2009a; Nakajima *et al.*, 2012). Although the reactions of the PAO/phyllobilin pathway have largely been elucidated during leaf senescence, there is a great body of evidence that points to its involvement also in other tissues and during adverse stress conditions. For example, natural green fruit and seed varieties are known from several species, such as tomato *green flesh*, bell pepper *chlorophyll retainer*, soybean (*Glycine max*) *d1d2*, and several green cotyledon varieties of pea. In each of these cases, *NYE* genes were shown to be mutated (Aubry *et al.*, 2008; Barry and Pandey, 2009; Fang *et al.*, 2014; Sato *et al.*, 2007). Furthermore, phyllobilins have been described

from ripening fruits of many species such as pear (*Pyrus communis*), apple (*Malus domestica*) and banana (Moser *et al.*, 2008a; Moser *et al.*, 2012; Müller *et al.*, 2007), and CCE proteins identified in fruit proteomics studies (Barsan *et al.*, 2010; Wang *et al.*, 2013). Likewise, an involvement of CCEs was shown for different stress conditions that induce degreening. For example, PAO induction and phyllobilin formation occurs during dehydration of resurrection plants (Christ *et al.*, 2014), and a role of PAO was shown for the hypersensitive response of *Arabidopsis* upon *Pseudomonas syringae* infection (Mur *et al.*, 2010).

All these data indicate high conservation of the PAO/phyllobilin pathway for Chl degradation under diverse conditions and in diverse plant tissues. Nevertheless, there are also some inconsistencies regarding Chl breakdown in leaves as compared to fruits and seeds. For example, tomato and *Arabidopsis* lines that are deficient in PPH exhibit a stay-green leaf phenotype, but fruit or seed degreening are unaffected (Guyer *et al.*, 2014; Zhang *et al.*, 2014). This hints to the involvement of additional, so far unknown, hydrolytic activities that are required for Chl breakdown in seeds and fruits. It has to be noted that in some instances, like during fruit ripening in *Citrus*, CLHs have been proposed to be involved in dephytylation (Azoulay-Shemer *et al.*, 2011; Azoulay Shemer *et al.*, 2008). At present, it can therefore not be excluded that CLHs might play a role in fruits. Similarly, while senescent leaves of the *Arabidopsis nye1-1* mutant that is deficient in NYE1 stay green during senescence, a block of Chl degradation during seed maturation is only evident in a NYE1/NYE2 double mutant (Delmas *et al.*, 2013; Wu *et al.*, 2016; Zhang *et al.*, 2014), pointing to functional redundancy in seeds that is less obvious in leaves. Also the part II CCEs of the PAO/phyllobilin pathway seem to be differentially active in different *Arabidopsis* tissues, as can be deduced from the relative abundancies of respective phyllobilins that show major differences between seeds and senescent leaves (Menghini and Hörtensteiner, unpublished).

Evolutionary aspects of the PAO/phyllobilin pathway

Apart from a few isolated and rather unclear reports of Chl degradation products in green algae (Doi *et al.*, 1997; Grabski *et al.*, 2016; Miyake *et al.*, 1995) and the well-studied example of *A. protothecoides*, which has been shown to degrade Chl to RCC-like products that are excreted (Engel *et al.*, 1996; Hörtensteiner *et al.*, 2000a; Oshio and Hase, 1969), phyllobilins have exclusively been identified in Angiosperms. This raises the intriguing question whether the PAO/phyllobilin pathway is conserved within all photosynthetic

organisms or whether Chl breakdown through this pathway evolved rather late and is (with maybe few exceptions) indeed restricted to Angiosperms. Phylogenetic analysis of NYEs, PPH, PAO, RCCR and TIC55 demonstrated that these enzymes are highly conserved within land plants, but sequence similarity drops significantly in lower photosynthetic organisms (Hauenstein *et al.*, 2016; Pružinská *et al.*, 2007; Schelbert *et al.*, 2009; Thomas *et al.*, 2009). Thus, it can be argued that all land plants may be capable to degrade Chl via the PAO/phyllobilin pathway and that the PAO/phyllobilin pathway evolved with the appearance of land plants (Hauenstein *et al.*, 2016). However, experimental support for this is scarce. RCCR homologs of the Gymnosperm *Pinus taeda*, the liverwort *Marchantia polymorpha* and the bryophyte *Physcomitrella patens* were shown to exhibit RCCR activity *in vitro* (Pružinská *et al.*, 2007). In line with the fact that *A. protothecoides* excretes RCC-like catabolites, RCCR is absent from green algae, and evolution of RCCR has been proposed a prerequisite for land colonization of terrestrial plants (Hörtensteiner, 2006).

Despite the presence of highly homologous CCE sequences within lower land plants, phyllobilins have so far not been identified in Gymnosperms, ferns or mosses. It was proposed that phyllobilins may not in all cases be the final products of Chl degradation. Accordingly, amounts of degraded Chl do not always correlate to the amounts of accumulated phyllobilins (Christ *et al.*, 2016; Losey and Engel, 2001) and monopyrrolic degradation products were proposed to occur in senescent barley leaves (Suzuki and Shioi, 1999; Suzuki *et al.*, 1999). It remains to be demonstrated to which extent degradation of Chl beyond the level of phyllobilins may occur, particularly in species like the primitive Gymnosperm *Ginkgo biloba* or the lycophyte *Selaginella moellendorffii* that are capable of leaf senescence but seem to be devoid of phyllobilins (Aubry & Hörtensteiner, unpublished).

Experimental evidence for the presence of CCE activities is also limited in green alga. The activity that produces RCC in *A. protothecoides* is biochemically rather similar to that of higher plants. Thus, both enzymes are monooxygenases that specifically incorporate a molecular oxygen-derived oxygen atom at the C1 formyl group (Curty *et al.*, 1995; Hörtensteiner *et al.*, 1998) and both activities are inhibited by Fe-chelators (Hörtensteiner *et al.*, 2000a; Hörtensteiner *et al.*, 1995). However, the most similar PAO-like enzyme encoded in the genome of *A. protothecoides* (Genbank accession number JAT76993) (Gao *et al.*, 2014), exhibits only 31% sequence identity with Arabidopsis PAO, while it is above 70% within land plants. It remains to be shown whether JAT76993 is a genuine PAO. Recently, an NYE

from *Chlamydomonas reinhardtii* was identified that exhibits Mg-dechelataase activity *in vitro* and after expression in Arabidopsis (Matsuda *et al.*, 2016).

The transcriptional regulation of the PAO/phyllobilin pathway

Since massive Chl breakdown is integral to green organ senescence, its regulation has frequently been investigated concomitantly with the general regulation of senescence. Upon consecutive cloning of *CCGs* during the last almost 20 years, it emerged that many *CCGs* are transcriptionally regulated during senescence. However, only during the past few years, the transcriptional regulation of *CCGs* was specifically explored, especially by yeast one-hybrid screening. This led to the recent identification of quite a few transcription regulators that directly target *CCGs* (Figure 2). Interestingly, almost all of these factors mediate the transcriptional regulation of both Chl breakdown and other senescence processes.

Involvement of EIN3, ORE1, and ERFs in directly activating major CCGs during ethylene-triggered Chl breakdown

The biosynthesis and signaling of ethylene (ET) have long been implicated in the process of degreening during leaf senescence and fruit ripening (Bleecker *et al.*, 1988; Chao *et al.*, 1997; Grbic and Bleecker, 1995; John *et al.*, 1995; Zacarias and Reid, 1990). Lines of solid evidence include: (i) the octuple Arabidopsis mutant plants of ACSs (*1-AMINOCYCLOPROPANE-1-CARBOXYLIC ACID SYNTHASES*), encoding the key enzyme catalyzing ethylene biosynthesis, is significantly delayed in degreening and senescence (Tsuchisaka *et al.*, 2009), (ii) EIN2 (ETHYLENE INSENSITIVE 2), the key upstream component of ethylene signaling, was found to upregulate *ORE1* (*ORESARAI*, *NAC092*), a key regulatory gene of degreening and leaf senescence, via negatively regulating *miR164* (*microRNA 164*) that directly targets the mRNA of *ORE1* (Kim *et al.*, 2009), and (iii) EIN3, the downstream target of EIN2, was shown to positively regulate both *ORE1* and *NAP* (*NAC-LIKE, ACTIVATED BY AP3/PI, NAC029*) directly (Kim *et al.*, 2014) or *ORE1* alone via inhibiting the expression of *miR164* (Li *et al.*, 2013). Although a few more NAC transcription factors were further identified downstream of *ORE1* (Kim *et al.*, 2014), it remained unclear until recently how ethylene signaling directly regulates Chl breakdown during leaf senescence. Qiu and colleagues (Qiu *et al.*, 2015) demonstrated that three major *CCGs*, *NYE1*, *NYC1*, and *PAO*, are the direct targets of EIN3 and *ORE1*, and, importantly, EIN3 and *ORE1* promote the transcription of *NYE1* and

NYC1 additively. Moreover, ORE1 promotes ethylene production by activating the expression of *ACS2*, a major ethylene biosynthesis gene. Therefore, EIN3, ORE1, and *ACS2* constitute a feed-forward loop involved in the regulation of ethylene-mediated Chl breakdown during leaf senescence in *Arabidopsis* (Qiu *et al.*, 2015). More recently, CitERF13, an ethylene responsive factor was identified in *Citrus sinensis*, which, like its homologs in *Arabidopsis* and tomato (*AtERF17* and *SlERF16*, respectively), acts to promote Chl breakdown, likely through activating the expression of *PPH* (Yin *et al.*, 2016). Above all, ethylene-triggered Chl breakdown is part of the degreening process during natural leaf senescence and fruit maturation.

Involvement of ABI3/ABI5/EEL, ABFs, and NAC016/029 in directly activating major CCGs during abscisic acid-triggered Chl breakdown

It has been repeatedly reported that the level of abscisic acid (ABA) is significantly or even dramatically increased during natural or stress-induced leaf senescence (Breeze *et al.*, 2011; Gepstein and Thimann, 1980; He *et al.*, 2005; Liu *et al.*, 2016). Furthermore, alterations in the endogenous ABA level or exogenous application of ABA or an ABA antagonist, significantly change the onset and/or progression of senescence, which is accompanied by changes in the expression profiles of *CCGs* (Raab *et al.*, 2009; Takasaki *et al.*, 2015; Yang *et al.*, 2014; Ye *et al.*, 2017). Notably, NAP, an important NAC transcription factor involved in the regulation of senescence, mediates ABA-triggered leaf senescence via NAP–SAG113 (a PROTEIN PHOSPHATASE 2C, PP2C) and NAP–AAO3 (ABSCISIC ALDEHYDE OXIDASE 3) regulatory modules in *Arabidopsis* (Yang *et al.*, 2014; Zhang and Gan, 2012). ABA-triggered leaf senescence has also been found to be mediated by RPK1 (RECEPTOR PROTEIN KINASE 1), MAPKKK18, SNAC-As (stress-responsive NAC transcription factors), NAC016, ABIG1 (ABA INSENSITIVE GROWTH 1), and an ABF (ABA-RESPONSIVE ELEMENT-BINDING FACTOR)/RAV1 (RELATED TO ABA-INSENSITIVE 3/VP1)–ORE1 regulatory module in *Arabidopsis* (Lee *et al.*, 2011; Liu *et al.*, 2016; Matsuoka *et al.*, 2015; Sakuraba *et al.*, 2016; Takasaki *et al.*, 2015; Zhao *et al.*, 2016).

Recently, the direct regulation of Chl breakdown by ABA signaling components has been explored in *Arabidopsis* as well as in rice. It was found that ABI5 (ABA INSENSITIVE 5) and EEL (ENHANCED EM LEVEL), two group A bZIP transcription factors that act downstream in the ABA signaling pathway, promote Chl breakdown by directly upregulating *NYE1* and *NYC1* (Sakuraba *et al.*, 2014a), and that NAC016, a newly identified component of

ABA signaling, also mediates the regulation of Chl breakdown *via* direct activation of *NYE1* (Sakuraba *et al.*, 2016). By screening for transcriptional regulators of *NYE1*, Gao *et al.* (2015) identified ABF2/3/4 as direct activators of *NYE1/2*, *NYC1*, and *PAO* in ABA-triggered as well as natural leaf senescence (Gao *et al.*, 2016). During seed maturation, ABI3, a B3 domain transcription factor, was shown to directly regulate the expression of *NYE1* and *NYE2* to promote Chl breakdown (Delmas *et al.*, 2013). In rice, OsNAP, an orthologue of AtNAP, was also identified as a key regulator of ABA-triggered and age-dependent leaf senescence, which directly targets *NYE1*, *NYC1*, *NYC3* (*PPH*), and *RCCR* as well as other *senescence-associated genes* (*SAGs*) (Liang *et al.*, 2014). This progress suggests that ABA-triggered Chl breakdown might be mainly responsible for degreening during abiotic stress-induced leaf senescence or during seed maturation.

Involvement of MYC2/3/4 and NAC019/055/072 in directly activating major CCGs during jasmonic acid-triggered Chl breakdown

Although quite a few decades ago, methyl jasmonate was shown to be a much stronger promoter of leaf senescence than ABA (Ueda and Kato, 1980), it was not until the beginning of this century that extensive evidence begun to accumulate for the involvement of endogenous jasmonic acid (JA) synthesis and signaling in regulating senescence (Breeze *et al.*, 2011; Castillo and Leon, 2008; He *et al.*, 2002; Lee *et al.*, 2015; Schommer *et al.*, 2008; Shan *et al.*, 2011; Yan *et al.*, 2012). Particularly, JA signaling repressors such as Rubisco activase and WRKY57 and antagonistic regulation of *SAG29* that involves the bHLH subgroup IIIe factors MYC2/3/4 and the bHLH subgroup IIId factors bHLH03/13/14/17 were recently identified to mediate JA-triggered leaf senescence (Jiang *et al.*, 2014; Qi *et al.*, 2015; Shan *et al.*, 2011). Strikingly, MYC2/3/4 were demonstrated to directly regulate *NYC1*, *NYE1* and *PAO* to promote Chl breakdown during JA-triggered leaf senescence (Zhu *et al.*, 2015) and NAC019/055/072, downstream targets of the MYCs in the JA signaling pathway, were also revealed to directly activate a similar set of *CCGs* (*NYE1/2*, *NYC1*) (Zhu *et al.*, 2015). In addition, MYC2 and NAC019 interact at the protein level to synergistically up-regulate the expression of *NYE1* (Zhu *et al.*, 2015). These findings suggest a complex hierarchical network of JA signaling in coordinating the regulation of efficient Chl breakdown, which may help to explain chlorosis incurred by pathogen infection.

Involvement of PIF4/5 and ABI5/EEL in directly activating and SOC1 in directly repressing

CCGs during dark-induced Chl breakdown

Light deprivation of green tissues (dark treatment) has been frequently exploited to investigate Chl breakdown and/or senescence processes. On the other hand, light has long been known to inhibit senescence of leaves *via* mediation of its perception and presumably signaling, and, specifically, phytochromes were suggested to be responsible for the senescence-inhibitory effect in light. This was rationalized from the observations that red light pulses were able to retard senescence almost as effective as white light, and the red light effect could be reverted by brief irradiation with far red light (Biswal and Biswal, 1984). The signaling details underlying have remained largely unknown for a long time; however, remarkable progress was made recently in elucidating the regulatory network involving phyB (phytochrome B) and PIFs (PHYTOCHROME-INTERACTING FACTORS) (Liebsch and Keech, 2016). It was shown that upon dark treatment *phyB* mutants exhibit an accelerated yellowing phenotype, whereas *phyB* overexpresser lines stayed green, indicating that the red light signal that inhibits senescence is mainly perceived by phyB (Sakuraba *et al.*, 2014a). Within the red light signaling pathway, PIF4/5 are major repressors downstream of phyB, which destabilizes them at the posttranslational level, while ELF3 (EARLY FLOWERING 3), a regulator of the circadian clock and flowering time, inhibits *PIF4/5* transcription (Leivar *et al.*, 2008; Nusinow *et al.*, 2011; Shin *et al.*, 2009). Notably, during dark-induced senescence, ABI5/EEL and EIN3 were revealed as downstream targets of PIF4/5, which, along with PIF4/5 themselves, regulate *ORE1* (Sakuraba *et al.*, 2014a). PIF4/5, as well as ABI5/EEL, EIN3, and ORE1, were independently demonstrated by different groups to directly regulate two key CCGs, *NYE1* and *NYC1* (Qiu *et al.*, 2015; Sakuraba *et al.*, 2014a; Song *et al.*, 2014; Zhang *et al.*, 2015). In addition, PIF4 was found to positively regulate the level of reactive oxygen species as well as ethylene biosynthesis and signaling, and to simultaneously repress the expression of *GLK* (*GOLDEN 2-LIKE*) genes, i.e. chloroplast activity maintenance genes (Oh *et al.*, 2012; Song *et al.*, 2014). These independent investigations collectively reveal PIF4 and PIF5, probably also PIF3, as the crucial players in the regulatory network of dark-/light deprivation-induced Chl breakdown and senescence, which hierarchically connects with known key regulators (Liebsch and Keech, 2016).

Nevertheless, the regulatory network of dark-induced senescence is far more complex than currently understood, as indicated by the fact that loss-of-functions of almost all SAGs results in stay-green phenotypes during dark-induced senescence. Furthermore, SOC1 (SUPPRESSOR OF OVEREXPRESSION OF CO 1), the key integrator of flowering

regulatory pathways, was recently identified as a negative regulator of dark-induced Chl breakdown/senescence (Chen *et al.*, 2017). SOC1 directly inhibits the expression of *NYE1* and *PPH* as well as of *SAG113*, which is involved in ABA-mediated leaf senescence (Zhang and Gan, 2012).

Conclusions

The ‘biological enigma’ of the fate of Chl (Hendry *et al.*, 1987) has been solved during the last three decades. We now know that a dedicated pathway, the PAO/phyllobilin pathway, is responsible for the degradation of Chl during leaf senescence and fruit ripening of plants. We know the end products of degradation, we know most of the genes and enzymes involved in the pathway, and gene regulatory aspects of the pathway are emerging. Nevertheless, several open questions on the PAO/phyllobilin pathway remain, for example, regarding the molecular identity of the activities that cause the formation of YCCs and PiCCs (Ulrich *et al.*, 2011; Vergeiner *et al.*, 2015) or that catalyze the widely-occurring C18-dihydroxylation (Kräutler *et al.*, 1991) (see Table 1). Furthermore, the fact that a single enzyme of a protein family, such as is the case for Arabidopsis MES16 and CYP89A9, has specifically been recruited to act in phyllobilin modification, favors that there might be a biological role(s) for phyllobilins, beyond their mere disposal as waste products inside the vacuole; however, such a role(s) has not been confirmed yet (Kräutler, 2016). Intriguingly, isolated reports have indicated a posttranscriptional regulation of PAO (Chung *et al.*, 2006; Pružinská *et al.*, 2003), but factors involved in such a regulatory mechanism are still missing. Finally, how all the CCGs are regulated so as to act in a coordinated manner also awaits to be elucidated.

Chl degradation through the PAO/phyllobilin pathway might not be the only way of degrading Chl. Indeed, in the past, peroxidases were repeatedly considered to be involved (Hynninen *et al.*, 2010; Johnson-Flanagan and Spencer, 1996; Matile, 1980); however, in none of the cases, have respective degradation products been identified. Recently, several reports (Kashiyama *et al.*, 2012; Kashiyama *et al.*, 2013) demonstrated Chl detoxification in heterotrophic protists during herbivory that does not involve porphyrin ring opening, but ring formation between the C17 propionic acid and ring E. These so-called 13²,17³-cyclopheophorbide enols are, in contrast to Chl, pheophytin and pheophorbide, nonfluorescent and incapable to produce singlet oxygen upon light excitation (Kashiyama *et al.*, 2012). Bioluminescent dinoflagellates and Euphausiids such as krill (*Euphausia pacifica*) use Chl-derived linear tetrapyrroles as

luciferins (Nakamura *et al.*, 1989; Nakamura *et al.*, 1988). Interestingly, in these cases the porphyrin ring is opened in the ‘Western’ position. The biochemical and molecular basis for the production of cyclopheophorbide enols or Chl-derived luciferins is unknown (Kashiyama and Tamiaki, 2014; Topalov and Kishi, 2001).

Acknowledgements

This work was supported by grants of the National Natural Science Foundation of China (31670287) to B.K., the National Postdoctoral Program for Innovative Talents (BX201700058) to J.C. and the Swiss National Science Foundation (31003A_149389/1 and 31CP30_163504/1) to S.H.

References

- Arkus KAJ, Cahoon EB, Jez JM. 2005. Mechanistic analysis of wheat chlorophyllase. Archives of Biochemistry and Biophysics **438**, 146-155.
- Armstead I, Donnison I, Aubry S, et al. 2006. From crop to model to crop: identifying the genetic basis of the staygreen mutation in the *Lolium/Festuca* forage and amenity grasses. New Phytologist **172**, 592-597.
- Armstead I, Donnison I, Aubry S, et al. 2007. Cross-species identification of Mendel's *I* locus. Science **315**, 73.
- Aubry S, Mani J, Hörtensteiner S. 2008. Stay-green protein, defective in Mendel's green cotyledon mutant, acts independent and upstream of pheophorbide *a* oxygenase in the chlorophyll catabolic pathway. Plant Molecular Biology **67**, 243-256.
- Azoulay-Shemer T, Harpaz-Saad S, Cohen-Peer R, et al. 2011. Dual N- and C-terminal processing of Citrus chlorophyllase precursor within the plastid membranes leads to the mature enzyme. Plant and Cell Physiology **52**, 70-83.
- Azoulay Shemer T, Harpaz-Saad S, Belausov E, Lovat N, Krokhin O, Spicer V, Standing KG, Goldschmidt EE, Eyal Y. 2008. Citrus chlorophyllase dynamics at ethylene-induced fruit color-break; a study of chlorophyllase expression, post-translational processing kinetics and *in-situ* intracellular localization. Plant Physiology **148**, 108-118.
- Balsera M, Soll J, Buchanan BB. 2010. Redox extends its regulatory reach to chloroplast protein import. Trends in Plant Science **15**, 515-521.
- Barry CS. 2009. The stay-green revolution: Recent progress in deciphering the mechanisms of chlorophyll degradation in higher plants. Plant Science **176**, 325-333.

- Barry CS, Pandey P.** 2009. A survey of cultivated heirloom tomato varieties identifies four new mutant alleles at the green-flesh locus. *Molecular Breeding* **24**, 269-276.
- Barsan C, Sanchez-Bel P, Rombaldi C, Egea I, Rossignol M, Kuntz M, Zouine M, Latche A, Bouzayen M, Pech JC.** 2010. Characteristics of the tomato chromoplast revealed by proteomic analysis. *Journal of Experimental Botany* **61**, 2413-2431.
- Berghold J, Breuker K, Oberhuber M, Hörtensteiner S, Kräutler B.** 2002. Chlorophyll breakdown in spinach: on the structure of five nonfluorescent chlorophyll catabolites. *Photosynthesis Research* **74**, 109-119.
- Berghold J, Eichmüller C, Hörtensteiner S, Kräutler B.** 2004. Chlorophyll breakdown in tobacco: on the structure of two nonfluorescent chlorophyll catabolites. *Chemistry & Biodiversity* **1**, 657-668.
- Berghold J, Müller T, Ulrich M, Hörtensteiner S, Kräutler B.** 2006. Chlorophyll breakdown in maize: on the structure of two nonfluorescent chlorophyll catabolites. *Monatshefte für Chemie* **137**, 751-763.
- Biswal UC, Biswal B.** 1984. Photocontrol of leaf senescence. *Photochemistry and Photobiology* **39**, 875-879.
- Bleecker AB, Estelle MA, Somerville C, Kende H.** 1988. Insensitivity to ethylene conferred by a dominant mutation in *Arabidopsis thaliana*. *Science* **241**, 1086-1089.
- Boij P, Patel R, Garcia C, Jarvis P, Aronsson H.** 2009. In vivo studies on the roles of Tic55-related proteins in chloroplast protein import in *Arabidopsis thaliana*. *Molecular Plant* **2**, 1397-1409.
- Breeze E, Harrison E, McHattie S, et al.** 2011. High-resolution temporal profiling of transcripts during *Arabidopsis* leaf senescence reveals a distinct chronology of processes and regulation. *The Plant Cell* **23**, 873-894.
- Büchert AM, Civello PM, Martínez GA.** 2011. Chlorophyllase versus pheophytinase as candidates for chlorophyll dephytylation during senescence of broccoli. *Journal of Plant Physiology* **168**, 337-343.
- Calibe A, Grimm R, Kaiser G, Lübeck J, Soll J, Heins L.** 1997. The chloroplastic protein import machinery contains a Rieske-type iron-sulfur cluster and a mononuclear iron-binding protein. *The EMBO Journal* **16**, 7342-7350.
- Castillo MC, Leon J.** 2008. Expression of the beta-oxidation gene 3-ketoacyl-CoA thiolase 2 (KAT2) is required for the timely onset of natural and dark-induced leaf senescence in *Arabidopsis*. *Journal of Experimental Botany* **59**, 2171-2179.
- Chao QM, Rothenberg M, Solano R, Roman G, Terzaghi W, Ecker JR.** 1997. Activation of the ethylene gas response pathway in *Arabidopsis* by the nuclear protein ETHYLENE-INSENSITIVE3 and related proteins. *Cell* **89**, 1133-1144.

- Chen J, Zhu X, Ren J, Qiu K, Li Z, Xie Z, Gao J, Zhou X, Kuai B.** 2017. Suppressor of overexpression of CO 1 negatively regulates dark-induced leaf degreening and senescence by directly repressing pheophytinase and other senescence-associated genes in *Arabidopsis*. *Plant Physiology* **173**, 1881-1891.
- Cheng YD, Guan JF.** 2014. Involvement of pheophytinase in ethylene-mediated chlorophyll degradation in the peel of harvested 'Yali' pear. *Journal of Plant Growth Regulation* **33**, 364-372.
- Christ B, Egert A, Süßenbacher I, Kräutler B, Bartels D, Peters S, Hörtensteiner S.** 2014. Water deficit induces chlorophyll degradation via the 'PAO/phyllobilin' pathway in leaves of homoio- (*Craterostigma pumilum*) and poikilochlorophyllous (*Xerophyta viscosa*) resurrection plants. *Plant Cell and Environment* **37**, 2521-2531.
- Christ B, Hauenstein M, Hörtensteiner S.** 2016. A liquid chromatography-mass spectrometry platform for the analysis of phyllobilins, the major degradation products of chlorophyll in *Arabidopsis thaliana*. *The Plant Journal* **88**, 505-518.
- Christ B, Hörtensteiner S.** 2014. Mechanism and significance of chlorophyll breakdown. *Journal of Plant Growth Regulation* **33**, 4-20.
- Christ B, Schelbert S, Aubry S, Süßenbacher I, Müller T, Kräutler B, Hörtensteiner S.** 2012. MES16, a member of the methylsterase protein family, specifically demethylates fluorescent chlorophyll catabolites during chlorophyll breakdown in *Arabidopsis*. *Plant Physiology* **158**, 628-641.
- Christ B, Süßenbacher I, Moser S, Bichsel N, Egert A, Müller T, Kräutler B, Hörtensteiner S.** 2013. Cytochrome P450 CYP89A9 is involved in the formation of major chlorophyll catabolites during leaf senescence in *Arabidopsis*. *The Plant Cell* **25**, 1868-1880.
- Chung DW, Pružinská A, Hörtensteiner S, Ort DR.** 2006. The role of pheophorbide *a* oxygenase expression and activity in the canola green seed problem. *Plant Physiology* **142**, 88-97.
- Costa ML, Civello PM, Chaves AR, Martinez GA.** 2002. Characterization of Mg-dechelataase activity obtained from *Fragaria x ananassa* fruit. *Plant Physiology and Biochemistry* **40**, 111-118.
- Curty C, Engel N.** 1996. Detection, isolation and structure elucidation of a chlorophyll *a* catabolite from autumnal senescent leaves of *Cercidiphyllum japonicum*. *Phytochemistry* **42**, 1531-1536.
- Curty C, Engel N, Gossauer A.** 1995. Evidence for a monooxygenase-catalyzed primary process in the catabolism of chlorophyll. *FEBS Letters* **364**, 41-44.
- Delmas F, Sankaranarayanan S, Deb S, Widdup E, Bournonville C, Bollier N, Northey JG, McCourt P, Samuel MA.** 2013. ABI3 controls embryo degreening through Mendel's I locus. *Proceedings of the National Academy of Sciences, USA* **110**, E3888-E3894.
- Djapic N, Djuric A, Pavlovic A.** 2009a. Chlorophyll biodegradation in *Vitis vinifera* var. *Pinot Noir* autumnal leaves. *Research Journal of Agricultural Science* **41**, 256-260.

- Djapic N, Pavlovic M.** 2008. Chlorophyll catabolite from *Parrotia persica* autumnal leaves. *Revista De Chimie* **59**, 878-882.
- Djapic N, Pavlovic M, Arsovski S, Vujic G.** 2009b. Chlorophyll biodegradation product from *Hamamelis virginiana* autumnal leaves. *Revista De Chimie* **60**, 398-402.
- Doi M, Shima S, Egashira T, Nakamura K, Okayama S.** 1997. New bile pigment excreted by a *Chlamydomonas reinhardtii* mutant: a possible breakdown catabolite of chlorophyll *a*. *Journal of Plant Physiology* **150**, 504-508.
- Drazkiewicz M.** 1994. Chlorophyllase: occurrence, functions, mechanism of action, effects of external and internal factors. *Photosynthetica* **30**, 321-331.
- Engel N, Curty C, Gossauer A.** 1996. Chlorophyll catabolism in *Chlorella protothecoides*. 8. Facts and artefacts. *Plant Physiology and Biochemistry* **34**, 77-83.
- Erhart T, Mittelberger C, Vergeiner C, Scherzer G, Holzner B, Robatscher P, Oberhuber M, Kräutler B.** 2016. Chlorophyll catabolites in senescent leaves of the plum tree (*Prunus domestica*). *Chemistry & Biodiversity* **13**, 1441-1453.
- Fang C, Li CC, Li WY, et al.** 2014. Concerted evolution of D1 and D2 to regulate chlorophyll degradation in soybean. *The Plant Journal* **77**, 700-712.
- Feller U, Anders I, Mae T.** 2008. Rubiscolytics: fate of Rubisco after its enzymatic function in a cell is terminated. *Journal of Experimental Botany* **59**, 1615-1624.
- Frankenberg N, Lagarias JC.** 2003. Biosynthesis and biological functions of bilins. In: Kadish KM, Smith KM, Guilard R, eds. *The Porphyrin Handbook*, Vol. 13/Chlorophylls and Bilins: Biosynthesis, Synthesis and Degradation. Amsterdam: Elsevier Science, 211-235.
- Frankenberg N, Mukougawa K, Kohchi T, Lagarias JC.** 2001. Functional genomic analysis of the HY2 family of ferredoxin-dependent bilin reductases from oxygenic photosynthetic organisms. *The Plant Cell* **13**, 965-978.
- Gao CF, Wang Y, Shen Y, Yan D, He X, Dai JB, Wu QY.** 2014. Oil accumulation mechanisms of the oleaginous microalga *Chlorella protothecoides* revealed through its genome, transcriptomes, and proteomes. *BMC Genomics* **15**.
- Gao S, Gao J, Zhu X, Song Y, Li Z, Ren G, Zhou X, Kuai B.** 2016. ABF2, ABF3, and ABF4 promote ABA-mediated chlorophyll degradation and leaf senescence by transcriptional activation of chlorophyll catabolic genes and senescence-associated genes in *Arabidopsis*. *Molecular Plant* **9**, 1272-1285.
- Gepstein S, Thimann KV.** 1980. Changes in the abscisic-acid content of oat leaves during senescence. *Proceedings of the National Academy of Sciences, USA* **77**, 2050-2053.
- Ginsburg S, Matile P.** 1993. Identification of catabolites of chlorophyll porphyrin in senescent rape cotyledons. *Plant Physiology* **102**, 521-527.

- Grabski K, Baranowski N, Skorko-Glonek J, Tukaj Z.** 2016. Chlorophyll catabolites in conditioned media of green microalga *Desmodesmus subspicatus*. *Journal of Applied Phycology* **28**, 889-896.
- Grbic V, Bleecker AB.** 1995. Ethylene regulates the timing of leaf senescence in *Arabidopsis*. *The Plant Journal* **8**, 595-602.
- Greenberg JT, Ausubel FM.** 1993. *Arabidopsis* mutants compromised for the control of cellular damage during pathogenesis and aging. *The Plant Journal* **4**, 327-341.
- Greenberg JT, Guo A, Klessig DF, Ausubel FM.** 1994. Programmed cell death in plants: a pathogen-triggered response activated coordinately with multiple defense functions. *Cell* **77**, 551-563.
- Guyer L, Schelbert Hofstetter S, Christ B, Silverstre Lira B, Rossi M, Hörtensteiner S.** 2014. Different mechanisms are responsible for chlorophyll dephytylation during fruit ripening and leaf senescence in tomato. *Plant Physiology* **166**, 44-56.
- Hauenstein M, Christ B, Das A, Aubry S, Hörtensteiner S.** 2016. A role for TIC55 as a hydroxylase of phyllobilins, the products of chlorophyll breakdown during plant senescence. *The Plant Cell* **28**, 2510-2527.
- He P, Osaki M, Takebe M, Shinano T, Wasaki J.** 2005. Endogenous hormones and expression of senescence-related genes in different senescent types of maize. *Journal of Experimental Botany* **56**, 1117-1128.
- He YH, Fukushige H, Hildebrand DF, Gan SS.** 2002. Evidence supporting a role of jasmonic acid in *Arabidopsis* leaf senescence. *Plant Physiology* **128**, 876-884.
- Hendry GAF, Houghton JD, Brown SB.** 1987. The degradation of chlorophyll - A biological enigma. *New Phytologist* **107**, 255-302.
- Hinder B, Schellenberg M, Rodoni S, Ginsburg S, Vogt E, Martinoia E, Matile P, Hörtensteiner S.** 1996. How plants dispose of chlorophyll catabolites. Directly energized uptake of tetrapyrrolic breakdown products into isolated vacuoles. *The Journal of Biological Chemistry* **271**, 27233-27236.
- Hirashima M, Tanaka R, Tanaka A.** 2009. Light-independent cell death induced by accumulation of pheophorbide *a* in *Arabidopsis thaliana*. *Plant and Cell Physiology* **50**, 719-729.
- Horie Y, Ito H, Kusaba M, Tanaka R, Tanaka A.** 2009. Participation of chlorophyll *b* reductase in the initial step of the degradation of light-harvesting chlorophyll *a/b*-protein complexes in *Arabidopsis*. *The Journal of Biological Chemistry* **284**, 17449-17456.
- Hörtensteiner S.** 2006. Chlorophyll degradation during senescence. *Annual Review of Plant Biology* **57**, 55-77.
- Hörtensteiner S.** 2009. Stay-green regulates chlorophyll and chlorophyll-binding protein degradation during senescence. *Trends in Plant Science* **14**, 155-162.

- Hörtensteiner S, Chinner J, Matile P, Thomas H, Donnison IS.** 2000a. Chlorophyll breakdown in *Chlorella protothecoides*: characterization of degreening and cloning of degreening-related genes. *Plant Molecular Biology* **42**, 439-450.
- Hörtensteiner S, Feller U.** 2002. Nitrogen metabolism and remobilization during senescence. *Journal of Experimental Botany* **53**, 927-937.
- Hörtensteiner S, Rodoni S, Schellenberg M, Vicentini F, Nandi OI, Qiu Y-L, Matile P.** 2000b. Evolution of chlorophyll degradation: the significance of RCC reductase. *Plant Biology* **2**, 63-67.
- Hörtensteiner S, Vicentini F, Matile P.** 1995. Chlorophyll breakdown in senescent cotyledons of rape, *Brassica napus* L.: enzymatic cleavage of pheophorbide *a in vitro*. *New Phytologist* **129**, 237-246.
- Hörtensteiner S, Wüthrich KL, Matile P, Ongania K-H, Kräutler B.** 1998. The key step in chlorophyll breakdown in higher plants. Cleavage of pheophorbide *a* macrocycle by a monooxygenase. *The Journal of Biological Chemistry* **273**, 15335-15339.
- Hu XY, Makita S, Schelbert S, Sano S, Ochiai M, Tsuchiya T, Hasegawa SF, Hörtensteiner S, Tanaka A, Tanaka R.** 2015. Reexamination of chlorophyllase function implies its involvement in defense against chewing herbivores. *Plant Physiology* **167**, 660-670.
- Hynninen PH, Kaartinen V, Kolehmainen E.** 2010. Horseradish peroxidase-catalyzed oxidation of chlorophyll *a* with hydrogen peroxide. Characterization of the products and mechanism of the reaction. *Biochimica et Biophysica Acta* **1797**, 531-542.
- Jakob-Wilk D, Holland D, Goldschmidt EE, Riov J, Eyal Y.** 1999. Chlorophyll breakdown by chlorophyllase: isolation and functional expression of the *Chlase1* gene from ethylene-treated *Citrus* fruit and its regulation during development. *The Plant Journal* **20**, 653-661.
- Jiang H, Li M, Liang N, Yan H, Wei Y, Xu X, Liu J, Xu Z, Chen F, Wu G.** 2007. Molecular cloning and function analysis of the *stay green* gene in rice. *The Plant Journal* **52**, 197-209.
- Jiang YJ, Liang G, Yang SZ, Yu DQ.** 2014. Arabidopsis WRKY57 functions as a node of convergence for jasmonic acid- and auxin-mediated signaling in jasmonic acid-induced leaf senescence. *The Plant Cell* **26**, 230-245.
- Jockusch S, Turro NJ, Banala S, Kräutler B.** 2014. Photochemical studies of a fluorescent chlorophyll catabolite - source of bright blue fluorescence in plant tissue and efficient sensitizer of singlet oxygen. *Photochemical & Photobiological Sciences* **13**, 407-411.
- John I, Drake R, Farrell A, Cooper W, Lee P, Horton P, Grierson D.** 1995. Delayed leaf senescence in ethylene-deficient ACC-oxidase antisense tomato plants - molecular and physiological analysis. *The Plant Journal* **7**, 483-490.

- Johnson-Flanagan AM, Spencer MS.** 1996. Chlorophyllase and peroxidase activity during degreening of maturing canola (*Brassica napus*) and mustard (*Brassica juncea*) seed. *Physiologia Plantarum* **97**, 353-359.
- Kashiyama Y, Tamiaki H.** 2014. Risk management by organisms of the phototoxicity of chlorophylls. *Chemistry Letters* **43**, 148-156.
- Kashiyama Y, Yokoyama A, Kinoshita Y, et al.** 2012. Ubiquity and quantitative significance of detoxification catabolism of chlorophyll associated with protistan herbivory. *Proceedings of the National Academy of Sciences, USA* **109**, 17328-17335.
- Kashiyama Y, Yokoyama A, Shiratori T, Inouye I, Kinoshita Y, Mizoguchi T, Tamiaki H.** 2013. $13^2,17^3$ -Cyclopheophorbide *b* enol as a catabolite of chlorophyll *b* in phycophagy by protists. *FEBS Letters* **587**, 2578-2583.
- Kim HJ, Hong SH, Kim YW, et al.** 2014. Gene regulatory cascade of senescence-associated NAC transcription factors activated by ETHYLENE-INSENSITIVE2-mediated leaf senescence signalling in *Arabidopsis*. *Journal of Experimental Botany* **65**, 4023-4036.
- Kim JH, Woo HR, Kim J, Lim PO, Lee IC, Choi SH, Hwang D, Nam HG.** 2009. Trifurcate feed-forward regulation of age-dependent cell death involving *miR164* in *Arabidopsis*. *Science* **323**, 1053-1057.
- Krätler B.** 2014. Phyllobilins - the abundant bilin-type tetrapyrrolic catabolites of the green plant pigment chlorophyll. *Chemical Society Reviews* **43**, 6227-6238.
- Krätler B.** 2016. Breakdown of chlorophyll in higher plants - Phyllobilins as abundant, yet hardly visible signs of ripening, senescence, and cell death. *Angewandte Chemie International Edition* **55**, 4882-4907.
- Krätler B, Hörtensteiner S.** 2014. Chlorophyll breakdown: chemistry, biochemistry and biology. In: Ferreira GC, Kadish KM, Smith KM, Guillard R, eds. *Handbook of Porphyrin Science - Chlorophyll, Photosynthesis and Bio-inspired Energy*, Vol. 28. Singapore: World Scientific Publishing, 117-185.
- Krätler B, Jaun B, Bortlik K-H, Schellenberg M, Matile P.** 1991. On the enigma of chlorophyll degradation: the constitution of a secoporphinoid catabolite. *Angewandte Chemie International Edition in English* **30**, 1315-1318.
- Kusaba M, Ito H, Morita R, et al.** 2007. Rice NON-YELLOW COLORING1 is involved in light-harvesting complex II and grana degradation during leaf senescence. *The Plant Cell* **19**, 1362-1375.
- Kusaba M, Tanaka A, Tanaka R.** 2013. Stay-green plants: what do they tell us about the molecular mechanism of leaf senescence. *Photosynthesis Research* **117**, 221-234.

- Lee IC, Hong SW, Whang SS, Lim PO, Nam HG, Koo JC.** 2011. Age-dependent action of an ABA-inducible receptor kinase, RPK1, as a positive regulator of senescence in *Arabidopsis* leaves. *Plant and Cell Physiology* **52**, 651-662.
- Lee SH, Sakuraba Y, Lee T, Kim KW, An G, Lee HY, Paek NC.** 2015. Mutation of *Oryza sativa* CORONATINE INSENSITIVE 1b (OsCOI1b) delays leaf senescence. *Journal of Integrative Plant Biology* **57**, 562-576.
- Leivar P, Monte E, Oka Y, Liu T, Carle C, Castillon A, Huq E, Quail PH.** 2008. Multiple phytochrome-interacting bHLH transcription factors repress premature seedling photomorphogenesis in darkness. *Current Biology* **18**, 1815-1823.
- Li HM, Pu H.** 2016. Crystal structure of methylesterase family member 16 (MES16) from *Arabidopsis thaliana*. *Biochemical and Biophysical Research Communications* **474**, 226-231.
- Li S, Gao J, Yao L, Ren G, Zhu X, Gao S, Qiu K, Zhou X, Kuai B.** 2016. The role of ANAC072 in the regulation of chlorophyll degradation during age- and dark-induced leaf senescence. *Plant Cell Reports* **35**, 1729-1741.
- Li ZH, Peng JY, Wen X, Guo HW.** 2013. ETHYLENE-INSENSITIVE3 is a senescence-associated gene that accelerates age-dependent leaf senescence by directly repressing miR164 transcription.
- Liang CZ, Wang YQ, Zhu YN, et al.** 2014. OsNAP connects abscisic acid and leaf senescence by fine-tuning abscisic acid biosynthesis and directly targeting senescence-associated genes in rice. *Proceedings of the National Academy of Sciences, USA* **111**, 10013-10018.
- Liao Y, An K, Zhou X, Chen W-J, Kuai B-K.** 2007. *AtCLH2*, a typical but possibly distinctive chlorophyllase gene in *Arabidopsis*. *Journal of Integrative Plant Biology* **49**, 531-539.
- Liebsch D, Keech O.** 2016. Dark-induced leaf senescence: new insights into a complex light-dependent regulatory pathway. *New Phytologist* **212**, 563-570.
- Lim PO, Kim HJ, Nam HG.** 2007. Leaf senescence. *Annual Review of Plant Biology* **58**, 115-136.
- Lin YP, Wu MC, Charng YY.** 2016. Identification of a chlorophyll dephytylase involved in chlorophyll turnover in *Arabidopsis*. *The Plant Cell* **28**, 2974-2990.
- Liu T, Longhurst AD, Talavera-Rauh F, Hokin SA, Barton MK.** 2016. The *Arabidopsis* transcription factor ABIG1 relays ABA signaled growth inhibition and drought induced senescence. *eLife* **5**, e13768.
- Losey FG, Engel N.** 2001. Isolation and characterization of a urobilinogenoidic chlorophyll catabolite from *Hordeum vulgare* L. *The Journal of Biological Chemistry* **276**, 27233-27236.
- Lu Y-P, Li Z-S, Drozdowicz Y-M, Hörtensteiner S, Martinoia E, Rea PA.** 1998. AtMRP2, an *Arabidopsis* ATP binding cassette transporter able to transport glutathione S-conjugates and chlorophyll catabolites: functional comparisons with AtMRP1. *The Plant Cell* **10**, 267-282.

- Mach JM, Castillo AR, Hoogstraten R, Greenberg JT.** 2001. The *Arabidopsis*-accelerated cell death gene *ACD2* encodes red chlorophyll catabolite reductase and suppresses the spread of disease symptoms. *Proceedings of the National Academy of Sciences, USA* **98**, 771-776.
- Makino A, Osmond B.** 1991. Effect of nitrogen nutrition on nitrogen partitioning between chloroplasts and mitochondria in pea and wheat. *Plant Physiology* **96**, 355-362.
- Matile P.** 1980. Catabolism of chlorophyll: involvement of peroxidase? *Zeitschrift für Pflanzenphysiologie* **99**, 457-478.
- Matile P.** 1997. The vacuole and cell senescence. In: Callow JA, ed. *Advances in Botanical Research*, Vol. 25. New York: Academic Press, 87-112.
- Matile P, Ginsburg S, Schellenberg M, Thomas H.** 1987. Catabolites of chlorophyll in senescent leaves. *Journal of Plant Physiology* **129**, 219-228.
- Matile P, Ginsburg S, Schellenberg M, Thomas H.** 1988. Catabolites of chlorophyll in senescing barley leaves are localized in the vacuoles of mesophyll cells. *Proceedings of the National Academy of Sciences, USA* **85**, 9529-9532.
- Matile P, Schellenberg M, Peisker C.** 1992. Production and release of a chlorophyll catabolite in isolated senescent chloroplasts. *Planta* **187**, 230-235.
- Matsuda K, Shimoda Y, Tanaka A, Ito H.** 2016. Chlorophyll *a* is a favorable substrate for *Chlamydomonas* Mg-dechelataase encoded by STAY-GREEN. *Plant Physiology and Biochemistry* **109**, 365-373.
- Matsuoka D, Yasufuku T, Furuya T, Nanmori T.** 2015. An abscisic acid inducible *Arabidopsis* MAPKKK, MAPKKK18 regulates leaf senescence via its kinase activity. *Plant Molecular Biology* **87**, 565-575.
- Meguro M, Ito H, Takabayashi A, Tanaka R, Tanaka A.** 2011. Identification of the 7-hydroxymethyl chlorophyll *a* reductase of the chlorophyll cycle in *Arabidopsis*. *The Plant Cell* **23**, 3442-3453.
- Mendel G.** 1866. Versuche über Pflanzenhybriden. *Verhandlungen des naturforschenden Vereines in Brünn* **4**, 3-47.
- Mittelberger C, Yalcinkaya H, Pichler C, et al.** 2017. Pathogen-induced leaf chlorosis: products of chlorophyll breakdown found in degreened leaves of Phytoplasma-infected apple (*Malus x domestica* Borkh.) and spricot (*Prunus armeniaca* L.) trees relate to the pheophorbide *a* oxygenase/phyllobilin pathway. *Journal of Agricultural and Food Chemistry* **65**, 2651-2660.
- Miyake K, Ohtomi M, Yoshizawa H, Sakamoto Y, Nakayama K, Okada M.** 1995. Water soluble pigments containing xylose and glucose in gametangia of the green alga, *Bryopsis maxima*. *Plant and Cell Physiology* **36**, 109-113.

- Morita R, Sato Y, Masuda Y, Nishimura M, Kusaba M.** 2009. Defect in non-yellow coloring 3, an a/b hydrolase-fold family protein, causes a stay-green phenotype during leaf senescence in rice. *The Plant Journal* **59**, 940-952.
- Moser S, Müller T, Ebert MO, Jockusch S, Turro NJ, Kräutler B.** 2008a. Blue luminescence of ripening bananas. *Angewandte Chemie International Edition* **47**, 8954-8957.
- Moser S, Müller T, Holzinger A, Lutz C, Jockusch S, Turro NJ, Kräutler B.** 2009a. Fluorescent chlorophyll catabolites in bananas light up blue halos of cell death. *Proceedings of the National Academy of Sciences, USA* **106**, 15538-15543.
- Moser S, Müller T, Holzinger A, Lutz C, Kräutler B.** 2012. Structures of chlorophyll catabolites in bananas (*Musa acuminata*) reveal a split path of chlorophyll breakdown in a ripening fruit. *Chemistry - A European Journal* **18**, 10873-10885.
- Moser S, Müller T, Oberhuber M, Kräutler B.** 2009b. Chlorophyll catabolites - Chemical and structural footprints of a fascinating biological phenomenon. *European Journal of Organic Chemistry* **2009**, 21-31.
- Moser S, Ulrich M, Müller T, Kräutler B.** 2008b. A yellow chlorophyll catabolite is a pigment of the fall colours. *Phytochemistry & Phytobiological Sciences* **7**, 1577-1581.
- Mühlecker W, Kräutler B.** 1996. Breakdown of chlorophyll: constitution of nonfluorescing chlorophyll-catabolites from senescent cotyledons of the dicot rape. *Plant Physiology and Biochemistry* **34**, 61-75.
- Mühlecker W, Kräutler B, Ginsburg S, Matile P.** 1993. Breakdown of chlorophyll: A tetrapyrrolic chlorophyll catabolite from senescent rape leaves. *Helvetica Chimica Acta* **76**, 2976-2980.
- Mühlecker W, Kräutler B, Moser D, Matile P, Hörtensteiner S.** 2000. Breakdown of chlorophyll: a fluorescent chlorophyll catabolite from sweet pepper (*Capsicum annuum*). *Helvetica Chimica Acta* **83**, 278-286.
- Mühlecker W, Ongania K-H, Kräutler B, Matile P, Hörtensteiner S.** 1997. Tracking down chlorophyll breakdown in plants: elucidation of the constitution of a 'fluorescent' chlorophyll catabolite. *Angewandte Chemie International Edition in English* **36**, 401-404.
- Müller T, Moser S, Ongania K-H, Pružinská A, Hörtensteiner S, Kräutler B.** 2006. A divergent path of chlorophyll breakdown in the model plant *Arabidopsis thaliana*. *ChemBioChem* **7**, 40-42.
- Müller T, Rafelsberger M, Vergeiner C, Kräutler B.** 2011. A dioxobilane as product of a divergent path of chlorophyll breakdown in Norway maple. *Angewandte Chemie International Edition* **50**, 10724-10727.
- Müller T, Ulrich M, Ongania KH, Kräutler B.** 2007. Colorless tetrapyrrolic chlorophyll catabolites found in ripening fruit are effective antioxidants. *Angewandte Chemie International Edition* **46**, 8699-8702.

- Müller T, Vergeiner S, Kräutler B.** 2014. Structure elucidation of chlorophyll catabolites (phyllobilins) by ESI-mass spectrometry - Pseudo-molecular ions and fragmentation analysis of a nonfluorescent chlorophyll catabolite (NCC). *International Journal of Mass Spectrometry* **365-366**, 48-55.
- Munné-Bosch S, Alegre L.** 2004. Die and let live: leaf senescence contributes to plant survival under drought stress. *Functional Plant Biology* **31**, 203-216.
- Mur LAJ, Aubry S, Mondhe M, et al.** 2010. Accumulation of chlorophyll catabolites photosensitizes the hypersensitive response elicited by *Pseudomonas syringae* in Arabidopsis. *New Phytologist* **188**, 161-174.
- Nakajima S, Ito H, Tanaka R, Tanaka A.** 2012. Chlorophyll *b* reductase plays an essential role in maturation and storability of Arabidopsis seeds. *Plant Physiology* **160**, 261-273.
- Nakamura H, Kishi Y, Shimomura O, Morse D, Hastings JW.** 1989. Structure of dinoflagellate luciferin and its enzymatic and nonenzymatic air-oxidation products. *Journal of the American Chemical Society* **111**, 7607-7611.
- Nakamura H, Musicki B, Kishi Y, Shimomura O.** 1988. Structure of the light emitter in krill (*Euphausia pacifica*) bioluminescence. *Journal of the American Chemical Society* **110**, 2683-2685.
- Nusinow DA, Helfer A, Hamilton EE, King JJ, Imaizumi T, Schultz TF, Farre EM, Kay SA.** 2011. The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* **475**, 398-402.
- Oberhuber M, Berghold J, Breuker K, Hörtensteiner S, Kräutler B.** 2003. Breakdown of chlorophyll: a nonenzymatic reaction accounts for the formation of the colorless "nonfluorescent" chlorophyll catabolites. *Proceedings of the National Academy of Sciences, USA* **100**, 6910-6915.
- Oberhuber M, Berghold J, Kräutler B.** 2008. Chlorophyll breakdown by a biomimetic route. *Angewandte Chemie International Edition* **47**, 3057-3061.
- Oberhuber M, Berghold J, Mühlecker W, Hörtensteiner S, Kräutler B.** 2001. Chlorophyll breakdown - on a nonfluorescent chlorophyll catabolite from spinach. *Helvetica Chimica Acta* **84**, 2615-2627.
- Oda-Yamamizo C, Mitsuda N, Sakamoto S, Ogawa D, Ohme-Takagi M, Ohmiya A.** 2016. The NAC transcription factor ANAC046 is a positive regulator of chlorophyll degradation and senescence in Arabidopsis leaves. *Scientific Reports* **6**, 23609.
- Oh E, Zhu JY, Wang ZY.** 2012. Interaction between BZR1 and PIF4 integrates brassinosteroid and environmental responses. *Nature Cell Biology* **14**, 802-809.

- Oshio Y, Hase E.** 1969. Studies on red pigments excreted by cells of *Chlorella protothecoides* during the process of bleaching induced by glucose or acetate. I. Chemical properties of the red pigments. *Plant and Cell Physiology* **10**, 41-49.
- Park S-Y, Yu J-W, Park J-S, et al.** 2007. The senescence-induced staygreen protein regulates chlorophyll degradation. *The Plant Cell* **19**, 1649-1664.
- Peoples MB, Dalling MJ.** 1988. The interplay between proteolysis and amino acid metabolism during senescence and nitrogen allocation. In: Noodén LD, Leopold AC, eds. *Senescence and Aging in Plants*. San Diego, USA: Academic Press, 181-217.
- Pružinská A, Anders I, Aubry S, Schenk N, Tapernoux-Lüthi E, Müller T, Kräutler B, Hörtensteiner S.** 2007. In vivo participation of red chlorophyll catabolite reductase in chlorophyll breakdown. *The Plant Cell* **19**, 369-387.
- Pružinská A, Anders I, Tanner G, Roca M, Hörtensteiner S.** 2003. Chlorophyll breakdown: pheophorbide *a* oxygenase is a Rieske-type iron-sulfur protein, encoded by the *accelerated cell death 1* gene. *Proceedings of the National Academy of Sciences, USA* **100**, 15259-15264.
- Pružinská A, Tanner G, Aubry S, et al.** 2005. Chlorophyll breakdown in senescent *Arabidopsis* leaves: characterization of chlorophyll catabolites and of chlorophyll catabolic enzymes involved in the degreening reaction. *Plant Physiology* **139**, 52-63.
- Qi TC, Wang JJ, Huang H, Liu B, Gao H, Liu YL, Song SS, Xie DX.** 2015. Regulation of jasmonate-induced leaf senescence by antagonism between bHLH subgroup IIIe and IIId factors in *Arabidopsis*. *The Plant Cell* **27**, 1634-1649.
- Qiu K, Li Z, Yang Z, et al.** 2015. EIN3 and ORE1 accelerate degreening during ethylene-mediated leaf senescence by directly activating chlorophyll catabolic genes in *Arabidopsis*. *PLoS Genetics* **11**, e1005399.
- Raab S, Drechsel G, Zarepour M, Hartung W, Koshiba T, Bittner F, Hoth S.** 2009. Identification of a novel E3 ubiquitin ligase that is required for suppression of premature senescence in *Arabidopsis*. *The Plant Journal* **59**, 39-51.
- Ren G, An K, Liao Y, Zhou X, Cao Y, Zhao H, Ge X, Kuai B.** 2007. Identification of a novel chloroplast protein AtNYE1 regulating chlorophyll degradation during leaf senescence in *Arabidopsis*. *Plant Physiology* **144**, 1429-1441.
- Ren GD, Zhou Q, Wu SX, Zhang YF, Zhang LG, Huang JR, Sun ZF, Kuai BK.** 2010. Reverse genetic identification of CRN1 and its distinctive role in chlorophyll degradation in *Arabidopsis*. *Journal of Integrative Plant Biology* **52**, 496-504.
- Rios JJ, Roca M, Perez-Galvez A.** 2014a. Nonfluorescent chlorophyll catabolites in loquat fruits (*Eriobotrya japonica* Lindl.). *Journal of Agricultural and Food Chemistry* **62**, 10576-10584.
- Rios JJ, Perez-Galvez A, Roca M.** 2014b. Non-fluorescent chlorophyll catabolites in quince fruits. *Food Research International* **65**, 255-262.

- Rios JJ, Roca M, Perez-Galvez A.** 2015. Systematic HPLC/ESI-high resolution-qTOF-MS methodology for metabolomic studies in nonfluorescent chlorophyll catabolites pathway. *Journal of Analytical Methods in Chemistry* **2015**, 490627.
- Rodoni S, Mühlecker W, Anderl M, Kräutler B, Moser D, Thomas H, Matile P, Hörtensteiner S.** 1997. Chlorophyll breakdown in senescent chloroplasts. Cleavage of pheophorbide *a* in two enzymic steps. *Plant Physiology* **115**, 669-676.
- Roiser MH, Müller T, Kräutler B.** 2015. Colorless chlorophyll catabolites in senescent florets of broccoli (*Brassica oleracea* var. *italica*). *Journal of Agricultural and Food Chemistry* **63**, 1385-1392.
- Saga Y, Tamiaki H.** 2012. Demetalation of chlorophyll pigments. *Chemistry & Biodiversity* **9**, 1659-1683.
- Sakuraba Y, Han SH, Lee SH, Hörtensteiner S, Paek NC.** 2016. *Arabidopsis* NAC016 promotes chlorophyll breakdown by directly upregulating *STAYGREEN1* transcription. *Plant Cell Reports* **35**, 155-166.
- Sakuraba Y, Jeong J, Kang MY, Kim J, Paek NC, Choi G.** 2014a. Phytochrome-interacting transcription factors PIF4 and PIF5 induce leaf senescence in *Arabidopsis*. *Nature Communications* **5**, doi: 10.1038/ncomms5636.
- Sakuraba Y, Kim D, Kim YS, Hörtensteiner S, Paek NC.** 2014b. *Arabidopsis* STAYGREEN-LIKE (SGRL) promotes abiotic stress-induced leaf yellowing during vegetative growth. *FEBS Letters* **588**, 3830-3837.
- Sakuraba Y, Park SY, Kim YS, Wang SH, Yoo SC, Hörtensteiner S, Paek NC.** 2014c. *Arabidopsis* STAY-GREEN2 is a negative regulator of chlorophyll degradation during leaf senescence. *Molecular Plant* **7**, 1288-1302.
- Sakuraba Y, Schelbert S, Park S-Y, Han S-H, Lee B-D, Besagni Andrès C, Kessler F, Hörtensteiner S, Paek N-C.** 2012. STAY-GREEN and chlorophyll catabolic enzymes interact at light-harvesting complex II for chlorophyll detoxification during leaf senescence in *Arabidopsis*. *The Plant Cell* **24**, 507-518.
- Sato Y, Moria R, Katsuma S, Nishimura M, Tanaka A, Kusaba M.** 2009. Two short-chain dehydrogenase/reductases, NON-YELLOW COLORING 1 and NYC1-LIKE, are required for chlorophyll *b* and light-harvesting complex II degradation during senescence in rice. *The Plant Journal* **57**, 120-131.
- Sato Y, Morita R, Nishimura M, Yamaguchi H, Kusaba M.** 2007. Mendel's green cotyledon gene encodes a positive regulator of the chlorophyll-degrading pathway. *Proceedings of the National Academy of Sciences, USA* **104**, 14169-14174.

- Schelbert S, Aubry S, Burla B, Agne B, Kessler F, Krupinska K, Hörtensteiner S.** 2009. Pheophytin pheophorbide hydrolase (pheophytinase) is involved in chlorophyll breakdown during leaf senescence in *Arabidopsis*. *The Plant Cell* **21**, 767-785.
- Schenk N, Schelbert S, Kanwischer M, Goldschmidt EE, Dörmann P, Hörtensteiner S.** 2007. The chlorophyllases AtCLH1 and AtCLH2 are not essential for senescence-related chlorophyll breakdown in *Arabidopsis thaliana*. *FEBS Letters* **581**, 5517-5525.
- Scherl M, Müller T, Kräutler B.** 2012. Chlorophyll catabolites in senescent leaves of the lime tree (*Tilia cordata*). *Chemistry & Biodiversity* **9**, 2605-2617.
- Scherl M, Müller T, Kreutz CR, Huber RG, Zass E, Liedl KR, Kräutler B.** 2016. Chlorophyll catabolites in fall leaves of the wych elm tree present a novel glycosylation motif. *Chemistry - A European Journal* **22**, 9498-9503.
- Schommer C, Palatnik JF, Aggarwal P, Chetelat A, Cubas P, Farmer EE, Nath U, Weigel D.** 2008. Control of jasmonate biosynthesis and senescence by miR319 targets. *PLoS Biology* **6**, 1991-2001.
- Schuler MA, Duan H, Bilgin M, Ali S.** 2006. *Arabidopsis* cytochrome P450s through the looking glass: a window on plant biochemistry. *Phytochemistry Reviews* **5**, 205-237.
- Shan XY, Wang JX, Chua LL, Jiang DA, Peng W, Xie DX.** 2011. The role of *Arabidopsis* Rubisco activase in jasmonate-induced leaf senescence. *Plant Physiology* **155**, 751-764.
- Shimoda Y, Ito H, Tanaka A.** 2012. Conversion of chlorophyll b to chlorophyll a precedes magnesium dechelation for protection against necrosis in *Arabidopsis*. *The Plant Journal* **72**, 501-511.
- Shimoda Y, Ito H, Tanaka A.** 2016. *Arabidopsis* STAY-GREEN, Mendel's green cotyledon gene, encodes magnesium-dechelataase. *The Plant Cell* **28**, 2147-2160.
- Shin J, Kim K, Kang H, Zulfugarov IS, Bae G, Lee CH, Lee D, Choi G.** 2009. Phytochromes promote seedling light responses by inhibiting four negatively-acting phytochrome-interacting factors. *Proceedings of the National Academy of Sciences, USA* **106**, 7660-7665.
- Shioi Y, Sasa T.** 1986. Purification of solubilized chlorophyllase from *Chlorella protothecoides*. *Methods in Enzymology* **123**, 421-427.
- Shioi Y, Watanabe K, Takamiya K.** 1996. Enzymatic conversion of pheophorbide *a* to a precursor of pyropheophorbide *a* in leaves of *Chenopodium album*. *Plant and Cell Physiology* **37**, 1143-1149.
- Song Y, Yang CW, Gao S, Zhang W, Li L, Kuai BK.** 2014. Age-triggered and dark-induced leaf senescence require the bHLH transcription factors PIF3, 4, and 5. *Molecular Plant* **7**, 1776-1787.
- Sugishima M, Kitamori Y, Noguchi M, Kohchi T, Fukuyama K.** 2009. Crystal structure of red chlorophyll catabolite reductase: enlargement of the ferredoxin-dependent bilin reductase family. *Journal of Molecular Biology* **389**, 376-387.

- Sugishima M, Okamoto Y, Noguchi M, Kohchi T, Tamiaki H, Fukuyama K.** 2010. Crystal structures of the substrate-bound forms of red chlorophyll catabolite reductase: implications for site-specific and stereospecific reaction. *Journal of Molecular Biology* **402**, 879-891.
- Süssenbacher I, Christ B, Hörtensteiner S, Kräutler B.** 2014. Hydroxymethylated phyllobilins: A puzzling new feature of the dioxobilin branch of chlorophyll breakdown. *Chemistry - A European Journal* **20**, 87-92.
- Süssenbacher I, Christ B, Hörtensteiner S, Kräutler B.** 2015. Hydroxymethylated phyllobilins in senescent *Arabidopsis thaliana* leaves - sign of a puzzling intermezzo of chlorophyll breakdown. *Chemistry - A European Journal* **21**, 11664-11670.
- Suzuki T, Shioi Y.** 2002. Re-examination of Mg-dechelation reaction in the degradation of chlorophylls using chlorophyllin *a* as substrate. *Photosynthesis Research* **74**, 217-223.
- Suzuki Y, Amano T, Shioi Y.** 2006. Characterization and cloning of the chlorophyll-degrading enzyme pheophorbidease from cotyledons of radish. *Plant Physiology* **140**, 716-725.
- Suzuki Y, Shioi Y.** 1999. Detection of chlorophyll breakdown products in the senescent leaves of higher plants. *Plant and Cell Physiology* **40**, 909-915.
- Suzuki Y, Tanabe K, Shioi Y.** 1999. Determination of chemical oxidation products of chlorophyll and porphyrin by high-performance liquid chromatography. *Journal of Chromatography A* **839**, 85-91.
- Takamiya K, Tsuchiya T, Ohta H.** 2000. Degradation pathway(s) of chlorophyll: what has gene cloning revealed? *Trends in Plant Science* **5**, 426-431.
- Takasaki H, Maruyama K, Takahashi F, Fujita M, Yoshida T, Nakashima K, Myouga F, Toyooka K, Yamaguchi-Shinozaki K, Shinozaki K.** 2015. SNAC-As, stress-responsive NAC transcription factors, mediate ABA-inducible leaf senescence. *The Plant Journal* **84**, 1114-1123.
- Tanaka R, Hirashima M, Satoh S, Tanaka A.** 2003. The *Arabidopsis-accelerated cell death* gene *ACD1* is involved in oxygenation of pheophorbide *a*: inhibition of pheophorbide *a* oxygenase activity does not lead to the "stay-green" phenotype in *Arabidopsis*. *Plant and Cell Physiology* **44**, 1266-1274.
- Tanaka R, Kobayashi K, Masuda T.** 2011. Tetrapyrrole metabolism in *Arabidopsis thaliana*. *The Arabidopsis Book* **9**, e0145.
- Tanaka R, Tanaka A.** 2011. Chlorophyll cycle regulates the construction and destruction of the light-harvesting complexes. *Biochimica et Biophysica Acta* **1807**, 968-976.
- Taylor L, Nunes-Nesi A, Parsley K, Leiss A, Leach G, Coates S, Wingler A, Fernie AR, Hibberd JM.** 2010. Cytosolic pyruvate, orthophosphate dikinase functions in nitrogen remobilization during leaf senescence and limits individual seed growth and nitrogen content. *The Plant Journal* **62**, 641-652.

- Thomas H, Huang L, Young M, Ougham H.** 2009. Evolution of plant senescence. *BMC Evolutionary Biology* **9**, 163.
- Tommasini R, Vogt E, Fromenteau M, Hörtensteiner S, Matile P, Amrhein N, Martinoia E.** 1998. An ABC transporter of *Arabidopsis thaliana* has both glutathione-conjugate and chlorophyll catabolite transport activity. *The Plant Journal* **13**, 773-780.
- Topalov G, Kishi Y.** 2001. Chlorophyll catabolism leading to the skeleton of dinoflagellate and krill luciferins: Hypothesis and model studies. *Angewandte Chemie International Edition* **40**, 3892-3894.
- Tsuchisaka A, Yu GX, Jin HL, Alonso JM, Ecker JR, Zhang XM, Gao S, Theologis A.** 2009. A combinatorial interplay among the 1-aminocyclopropane-1-carboxylate isoforms regulates ethylene biosynthesis in *Arabidopsis thaliana*. *Genetics* **183**, 979-1003.
- Tsuchiya T, Ohta H, Okawa K, Iwamatsu A, Shimada H, Masuda T, Takamiya K.** 1999. Cloning of chlorophyllase, the key enzyme in chlorophyll degradation: finding of a lipase motif and the induction by methyl jasmonate. *Proceedings of the National Academy of Sciences, USA* **96**, 15362-15367.
- Tu SL, Gunn A, Toney MD, Britt RD, Lagarias JC.** 2004. Biliverdin reduction by cyanobacterial phycocyanobilin : ferredoxin oxidoreductase (PcyA) proceeds via linear tetrapyrrole radical intermediates. *Journal of the American Chemical Society* **126**, 8682-8693.
- Tuba Z, Lichtenthaler HK, Csintalan Z, Nagy Z, Sente K.** 1996. Loss of chlorophylls, cessation of photosynthetic CO₂ assimilation and respiration in the poikilochlorophyllous plant *Xerophyta scabrata* during desiccation. *Physiologia Plantarum* **96**, 383-388.
- Ueda J, Kato J.** 1980. Isolation and identification of a senescence-promoting substance from wormwood (*Artemisia absinthium* L.). *Plant Physiology* **66**, 246-249.
- Ulrich M, Moser S, Müller T, Kräutler B.** 2011. How the colourless 'nonfluorescent' chlorophyll catabolites rust. *Chemistry - A European Journal* **17**, 2330-2334.
- Unno M, Matsui T, Ikeda-Saito M.** 2007. Structure and catalytic mechanism of heme oxygenase. *Natural Product Reports* **24**, 553-570.
- Vergara-Dominguez H, Rios JJ, Gandul-Rojas B, Roca M.** 2016. Chlorophyll catabolism in olive fruits (var. Arbequina and Hojiblanca) during maturation. *Food Chemistry* **212**, 604-611.
- Vergeiner C, Ulrich M, Li CJ, Liu XJ, Müller T, Kräutler B.** 2015. Stereo- and regioselective phyllobilane oxidation in leaf homogenates of the peace lily (*Spathiphyllum wallisii*): hypothetical endogenous path to yellow chlorophyll catabolites. *Chemistry - A European Journal* **21**, 136-149.
- Vicentini F, Iten F, Matile P.** 1995. Development of an assay for Mg-dechelataase of oilseed rape cotyledons, using chlorophyllin as the substrate. *Physiologia Plantarum* **94**, 57-63.

- vom Dorp K, Hölzl G, Plohm C, Eisenhut M, Abraham M, Weber APM, Hanson AD, Dörmann P.** 2015. Remobilization of phytol from chlorophyll degradation is essential for tocopherol synthesis and growth of *Arabidopsis*. *The Plant Cell* **27**, 2846-2859.
- Wang YQ, Yang Y, Fei ZJ, Yuan H, Fish T, Thannhauser TW, Mazourek M, Kochian LV, Wang XW, Li L.** 2013. Proteomic analysis of chromoplasts from six crop species reveals insights into chromoplast function and development. *Journal of Experimental Botany* **64**, 949-961.
- Willstätter R, Stoll A.** 1913. Die Wirkungen der Chlorophyllase. In: Willstätter R, Stoll A, eds. *Untersuchungen über Chlorophyll*. Berlin: Verlag Julius Springer, 172-187.
- Wu S, Li Z, Yang L, et al.** 2016. NON-YELLOWING2 (NYE2), a close paralog of NYE1, plays a positive role in chlorophyll degradation in *Arabidopsis*. *Molecular Plant* **9**, 624-627.
- Wüthrich KL, Bovet L, Hunziker PE, Donnison IS, Hörtensteiner S.** 2000. Molecular cloning, functional expression and characterisation of RCC reductase involved in chlorophyll catabolism. *The Plant Journal* **21**, 189-198.
- Xodo LE, Rapozzi V, Zacchigna M, Drioli S, Zorzet S.** 2012. The chlorophyll catabolite pheophorbide *a* as a photosensitizer for the photodynamic therapy. *Current Medicinal Chemistry* **19**, 799-807.
- Yan YX, Christensen S, Isakeit T, Engelberth J, Meeley R, Hayward A, Emery RJN, Kolomiets MV.** 2012. Disruption of OPR7 and OPR8 reveals the versatile functions of jasmonic acid in maize development and defense. *The Plant Cell* **24**, 1420-1436.
- Yang JD, Worley E, Udvardi M.** 2014. A NAP-AAO3 regulatory module promotes chlorophyll degradation via ABA biosynthesis in *Arabidopsis* leaves. *The Plant Cell* **26**, 4862-4874.
- Yang Y, Xu R, Ma CJ, Vlot AC, Klessig DF, Pichersky E.** 2008. Inactive methyl indole-3-acetic acid ester can be hydrolyzed and activated by several esterases belonging to the *AtMES* esterase family of *Arabidopsis*. *Plant Physiology* **147**, 1034-1045.
- Ye Y, Zhou L, Liu X, Liu H, Li D, Cao M, Chen H, Xu L, Zhu JK, Zhao Y.** 2017. A novel chemical inhibitor of ABA signaling targets all ABA receptors. *Plant Physiology* **173**, 2356-2369.
- Yin XR, Xie XL, Xia XJ, Yu JQ, Ferguson IB, Giovannoni JJ, Chen KS.** 2016. Involvement of an ethylene response factor in chlorophyll degradation during citrus fruit degreening. *The Plant Journal* **86**, 403-412.
- Zacarias L, Reid MS.** 1990. Role of growth-regulators in the senescence of *Arabidopsis thaliana* leaves. *Physiologia Plantarum* **80**, 549-554.
- Zhang J, Yu G, Wen W, Ma X, Xu B, Huang B.** 2016. Functional characterization and hormonal regulation of the PHEOPHYTINASE gene *LpPPH* controlling leaf senescence in perennial ryegrass. *Journal of Experimental Botany* **67**, 935-945.

- Zhang KW, Gan SS.** 2012. An abscisic acid-AtNAP transcription factor-SAG113 protein phosphatase 2C regulatory chain for controlling dehydration in senescing Arabidopsis leaves. *Plant Physiology* **158**, 961-969.
- Zhang W, Liu TQ, Ren GD, Hörtensteiner S, Zhou YM, Cahoon EB, Zhang CY.** 2014. Chlorophyll degradation: the tocopherol biosynthesis related phytol hydrolase in Arabidopsis seeds is still missing. *Plant Physiology* **166**, 70-79.
- Zhang YQ, Liu ZJ, Chen YD, He JX, Bi YR.** 2015. PHYTOCHROME-INTERACTING FACTOR 5 (PIF5) positively regulates dark-induced senescence and chlorophyll degradation in Arabidopsis. *Plant Science* **237**, 57-68.
- Zhao Y, Chan ZL, Gao JH, et al.** 2016. ABA receptor PYL9 promotes drought resistance and leaf senescence. *Proceedings of the National Academy of Sciences, USA* **113**, 1949-1954.
- Zhou X, Liao Y, Ren GD, Zhang YY, Chen WJ, Kuai BK.** 2007. Repression of AtCLH1 expression results in a decrease in the ratio of chlorophyll *a/b* but does not affect the rate of chlorophyll degradation during leaf senescence. *Journal of Plant Physiology and Molecular Biology* **33**, 596-606.
- Zhu XY, Chen JY, Xie ZK, Gao J, Ren GD, Gao S, Zhou X, Kuai BK.** 2015. Jasmonic acid promotes degreening via MYC2/3/4-and ANAC019/055/072-mediated regulation of major chlorophyll catabolic genes. *The Plant Journal* **84**, 597-610.

Table 1. Nonfluorescent phytylobilins identified to date.

Classifying Name ^a	m/z [M+H] ⁺ calculated	Formula (M)	Side chain modifications ^b					Members of phytylobilin class	Reference ^c
			R ¹ (C3 ²)	R ² (O8 ⁴)	R ³ (C18)	R ⁴ (C2/C4)	R ⁵ (C12 ³)		
DNCC_602	603.2813	C ₃₃ H ₃₈ N ₄ O ₇	H	H	vinyl	H	H	<i>At</i> -DNCC-45 ⁱ , <i>At</i> -DNCC-48 ⁱ	[1]
DNCC_616	617.2970	C ₃₄ H ₄₀ N ₄ O ₇	H	CH ₃	vinyl	H	H	<i>At-mes16</i> -DNCC-47	[2]
DNCC_618	619.2762	C ₃₃ H ₃₈ N ₄ O ₈	OH	H	vinyl	H	H	<i>At</i> -DNCC-1, <i>At</i> -DNCC-33; <i>Bo</i> -DNCC	[1,3,4]
DNCC_632-1	633.2919	C ₃₄ H ₄₀ N ₄ O ₈	H	H	vinyl	C4-HM	H	<i>At</i> -4HM-DNCC-41	[1]
DNCC_632-2	633.2919	C ₃₄ H ₄₀ N ₄ O ₈	H	H	vinyl	C2-HM	H	<i>At</i> -2HM- <i>iso</i> -DNCC-42	[1]
DNCC_632-3	633.2919	C ₃₄ H ₄₀ N ₄ O ₈	OH	CH ₃	vinyl	H	H	<i>At-mes16</i> -DNCC-38, <i>Hvir</i> -UNCC, <i>Pa</i> -DNCC-53, <i>Pp</i> -UNCC, <i>Vv</i> -UNCC	[2,5-7,31]
DNCC_646-1	647.3075	C ₃₅ H ₄₂ N ₄ O ₈	H	CH ₃	vinyl	C4-HM	H	<i>At-mes16</i> -9HM-DNCC-44	[2]
DNCC_646-2	647.3075	C ₃₅ H ₄₂ N ₄ O ₈	H	CH ₃	vinyl	C2-HM	H	<i>At-mes16</i> -7HM- <i>iso</i> -DNCC-46	[2]
DNCC_650 ^d	651.3030	C ₃₄ H ₄₂ N ₄ O ₉	H	CH ₃	DHethyl	H	H	-	[8]
DNCC_666	667.2979	C ₃₄ H ₄₂ N ₄ O ₁₀	OH	CH ₃	DHethyl	H	H	<i>Ap</i> -DNCC ^l , <i>Hv</i> -UCC ^k , <i>Co</i> -NDCC-2	[9,10,27]
DNCC_780	781.3291	C ₃₉ H ₄₈ N ₄ O ₁₃	OGlc	H	vinyl	H	H	-	[11]
DNCC_794	795.3447	C ₄₀ H ₅₀ N ₄ O ₁₃	OGlc	CH ₃	vinyl	H	H	-	[11]
DNCC_828 ^d	829.3507	C ₄₀ H ₅₂ N ₄ O ₁₅	OGlc	CH ₃	DHethyl	H	H	<i>Co</i> -NDCC-1	[27]
DNCC_880 ^e	881.3457	C ₄₃ H ₅₂ N ₄ O ₁₆	OGlcMal	CH ₃	vinyl	H	H	-	[8]
NCC_614	615.2813	C ₃₄ H ₃₈ N ₄ O ₇	H	H	vinyl	H	H	<i>At</i> -NCC-5, <i>Bn</i> -NCC-4, <i>Bo</i> -NCC-2, <i>Oe</i> -NCC-4	[4,12,13,32]
NCC_628	629.2970	C ₃₅ H ₄₀ N ₄ O ₇	H	CH ₃	vinyl	H	H	<i>Cj</i> -NCC-2, <i>Md</i> -NCC-58, <i>Oe</i> -NCC-3, <i>Pa</i> -NCC-58, <i>Pd</i> -NCC-71, <i>So</i> -NCC-5	[14-16,31,32]
NCC_630 ^f	631.2762	C ₃₄ H ₃₈ N ₄ O ₈	OH	H	vinyl	H	H	<i>At</i> -NCC-2, <i>Bn</i> -NCC-3, <i>Ej</i> -NCC-3, <i>Mc</i> -NCC-49, <i>Oe</i> -NCC-1, <i>So</i> -NCC-3	[12,16-19,32]
NCC_644	645.2919	C ₃₅ H ₄₀ N ₄ O ₈	OH	CH ₃	vinyl	H	H	<i>Cj</i> -NCC-1, <i>Ej</i> -NCC-4, <i>Mc</i> -NCC-61, <i>Md</i> -NCC-2, <i>Md</i> -NCC-49 epimer, <i>Md</i> -NCC-50, <i>Oe</i> -NCC-2, <i>Pa</i> -NCC-49, <i>Pa</i> -NCC-50, <i>Pc</i> -NCC-2, <i>Pd</i> -NCC-60, <i>So</i> -NCC-4, <i>Sw</i> -NCC-58	[14,16,18-22,31,32]
NCC_662 ^d	663.3030	C ₃₅ H ₄₂ N ₄ O ₉	H	CH ₃	DHethyl	H	H	-	[8]
NCC_664	665.2823	C ₃₄ H ₄₀ N ₄ O ₁₀	OH	H	DHethyl	H	H	<i>Mc</i> -NCC-26, <i>So</i> -NCC-1	[16,18]
NCC_678	679.2979	C ₃₅ H ₄₂ N ₄ O ₁₀	OH	CH ₃	DHethyl	H	H	<i>Hv</i> -NCC-1, <i>Ej</i> -NCC-1, <i>Mc</i> -NCC-42, <i>Md</i> -NCC-35, <i>Pa</i> -NCC-35, <i>Pd</i> -NCC-40, <i>So</i> -NCC-2	[14,18,19,23,24,31]
NCC_716	717.2772	C ₃₇ H ₄₀ N ₄ O ₁₁	OMal	H	vinyl	H	H	<i>Bn</i> -NCC-1	[25]
NCC_730	731.2928	C ₃₈ H ₄₂ N ₄ O ₁₁	OMal	CH ₃	vinyl	H	H	<i>Ej</i> -NCC-2	[19]
NCC_788	789.3347	C ₄₁ H ₄₈ N ₄ O ₁₂	OGlc ^h	CH ₃	vinyl	H	H ⁱ	<i>Ug</i> -NCC-53	[26]
NCC_792	793.3291	C ₄₀ H ₄₈ N ₄ O ₁₃	OGlc	H	vinyl	H	H	<i>At</i> -NCC-1, <i>Bn</i> -NCC-2, <i>Bo</i> -NCC-1	[4,12,17]
NCC_806	807.3447	C ₄₁ H ₅₀ N ₄ O ₁₃	OGlc	CH ₃	vinyl	H	H	<i>At</i> -NCC-4, <i>Mc</i> -NCC-59, <i>Md</i> -NCC-1, <i>Md</i> -NCC-47, <i>Nr</i> -NCC-2, <i>Pa</i> -NCC-47, <i>Pc</i> -NCC-1, <i>Pd</i> -NCC-56, <i>Tc</i> -NCC-2, <i>Ug</i> -NCC-43, <i>Zm</i> -NCC-2	[12,14,18,21,26,28-31]
NCC_826	827.3351	C ₄₀ H ₅₀ N ₄ O ₁₅	OGlc	H	DHethyl	H	H	<i>Co</i> -NCC-2	[27]
NCC_830	831.3089	C ₄₂ H ₄₆ N ₄ O ₁₄	OH	CH ₃	vinyl	H	daucyl	<i>Mc</i> -NCC-55 ^k , <i>Mc</i> -NCC-58 ^k	[18]
NCC_840	841.3507	C ₄₁ H ₅₂ N ₄ O ₁₅	OGlc	CH ₃	DHethyl	H	H	<i>Co</i> -NCC-1, <i>Md</i> -NCC-31, <i>Pa</i> -NCC-31, <i>Pa</i> -NCC-33, <i>Pd</i> -NCC-35, <i>Tc</i> -NCC-1, <i>Ug</i> -NCC-27, <i>Zm</i> -NCC-1	[14,26,27,29-31]
NCC_892	893.3457	C ₄₄ H ₅₂ N ₄ O ₁₆	OGlcMal	CH ₃	vinyl	H	H	<i>Nr</i> -NCC-1	[28]
NCC_926 ^g	927.3511	C ₄₄ H ₅₄ N ₄ O ₁₈	OGlcMal	CH ₃	DHethyl	H	H	-	[8]
NCC_1002	1003.404	C ₄₇ H ₆₂ N ₄ O ₂₀	OGlc	CH ₃	OH-OGlc	H	H	<i>Md</i> -NCC-29, <i>Pa</i> -NCC-29, <i>Pd</i> -NCC-32	[14,31]



^aclassifying nomenclature for phyllobilins as proposed by Christ *et al.* (2016). Note that this classification ignores isomeric properties of respective phyllobilin members. ^bpositions of modifications as depicted in Figure 1; DHethyl, -CH(OH)-CH₂OH; HM, -CH₂OH; Glc, glucosyl; Mal, malonyl; OH-OGlc, -CH(OH)-CH₂OGlc; ^creferences: [1] Süßenbacher *et al.* 2015, [2] Süßenbacher *et al.* 2014, [3] Christ *et al.* 2013, [4] Roiser *et al.* 2015, [5] Djapic & Pavlovic 2008, [6] Djapic *et al.* 2009a, [7] Djapic *et al.* 2009b, [8] Das & Hörtensteiner, unpublished, [9] Losey & Engel 2001, [10] Müller *et al.* 2011, [11] Christ *et al.* 2016, [12] Pruzinska *et al.* 2005, [13] Kräutler & Hörtensteiner 2014, [14] Erhart *et al.* 2016, [15] Oberhuber *et al.* 2003, [16] Berghold *et al.* 2002, [17] Mühlecker *et al.* 1996, [18] Moser *et al.* 2012, [19] Rios *et al.* 2014a, [20] Curty and Engel 1996, [21] Müller *et al.* 2007, [22] Vergeiner *et al.* 2015, [23] Kräutler *et al.* 1991, [24] Oberhuber *et al.* 2001, [25] Mühlecker *et al.* 1993, [26] Scherl *et al.* 2016, [27] Rios *et al.* 2014b, [28] Berghold *et al.* 2004, [29] Berghold *et al.* 2006, [30] Scherl *et al.* 2012, [31] Mittelberger *et al.* 2017, [32] Vergara-Dominguez *et al.* 2016; ^dcatabolite identified in barley; ^ecatabolite identified in *Lolium perenne*; ^fNCC_630 also for *At*-NCC-3, but there OH is at C2 instead of C3² (Müller *et al.* 2006); ^gcatabolite identified in barley, wheat (*Triticum aestivum*) and *Sorghum bicolor*; ^ha glucose moiety intramolecularly connects C3² and C12³; ⁱpotential C4 diastereomers; ^jC10 epimers; ^ktwo C4 diastereomers of *Hv*-UCC were identified.

Figure legends

Figure 1. The PAO/phyllobilin pathway of chlorophyll breakdown in leaves. **(A)** Reactions of the first part of the pathway from chlorophyll to hydroxy-*p*FCC that take place inside the chloroplast. **(B)** *p*FCC/hydroxy-*p*FCC-modifying reactions of the second part of the pathway that take place outside the chloroplast and lead to the diversity of phyllobilins. For porphyrins and phyllobilins, respectively, pyrrole rings (A-D) and relevant atoms are shown in chlorophyll and *p*FCC, respectively. R¹ to R⁵ indicate side chain modifications according to Table 1. See the text for abbreviations of enzyme names. Note, that MES16 and CYP89A9 are specific *Arabidopsis* enzymes, while all other enzymes are known from different plant species. Also note, that R⁴ in DFCCs and DNCCs can alternatively be positioned at C2 (see also Table 1). DFCCs, dioxobilin-type fluorescent chlorophyll catabolites; DNCCs, dioxobilin-type nonfluorescent chlorophyll catabolites; FCCs, formylxobilin-type fluorescent chlorophyll catabolites; NCCs, formylxobilin-type nonfluorescent chlorophyll catabolites; *p*FCC, *primary* fluorescent chlorophyll catabolite, RCC, red chlorophyll catabolite.

Figure 2. The up-to-date regulatory network of chlorophyll (Chl) breakdown. **(A)** Regulatory pathways of *Chl catabolic genes (CCGs)* by light and respective hormones. Arrows and lines ending in bars indicate promotions and suppressions, respectively; Solid lines represent solid evidence while a dotted line indicates indirect evidence. **(B)** Diagrams showing the known promoters, as well as a repressor in two cases, of individual *CCGs*. See the text for abbreviations and additional details.

A



B

